

Werk

Titel: Mathematische Annalen

Verlag: Springer

Jahr: 1989

Kollektion: Mathematica

Werk Id: PPN235181684_0283

PURL: http://resolver.sub.uni-goettingen.de/purl?PID=PPN235181684_0283 | LOG_0047

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

A Root System for the Lyons Group

Wolfram Neutsch¹ and Werner Meyer²

¹ Institut für Astrophysik der Universität Bonn, Auf dem Hügel 71, D-5300 Bonn 1, Federal Republic of Germany

² Max-Planck-Institut für Mathematik, Gottfried-Claren-Strasse 26, D-5300 Bonn 3, Federal Republic of Germany

0. Introduction

Sims (1973) proved the existence and uniqueness of the sporadic group Ly predicted by Lyons (1972) through the computer-aided construction of a presentation which, unfortunately, is rather cumbersome and does not lead to an insight into the structure of the group.

Meanwhile, much more information on Ly has become available.

Kantor (1981) found a Tits geometry for Ly which is “almost” a building. Meyer et al. (1985) gave the absolute minimal representation of Ly (111-dimensional over \mathbb{F}_3) which had been conjectured by Meyer and Neutsch (1984) and, independently, by Woldar (1987). Later, Wilson [1984, 1985] compiled the list of all maximal subgroups in Ly . His investigation uses the minimal representation explicitly, while the verification of the latter depends on Sims’ presentation. For that reason, it would be of great interest to have a simpler existence and uniqueness proof.

Inspired by Kantor’s results, we were led to the idea of giving a symmetric presentation of a group Γ by making use of the beautiful geometry of Ly .

Our relations are shown to be fulfilled by certain generators (“roots”) of the Lyons group, and most probably they define Ly itself.

The construction is carried out in a fashion nearly identical to the methods of Chevalley theory employed to study the Tits buildings of the groups of Lie type ($G_2(5) < Ly$ should be considered as a prototype). The geometric spirit of our presentation renders this possible. It is a first step towards an understanding of the Lyons group.

1. Relations in a Group of Ly Type

We say a group A is of Ly type if it has the following properties:

- (1) A is simple;
- (2) A contains an involution z with $C_A(z) \cong 2 \wedge A_{11}$.

Lyons [1972] shows that a group fulfilling (1) and (2) is of order

$$|Ly| = 2^8 \cdot 3^7 \cdot 5^6 \cdot 7 \cdot 11 \cdot 31 \cdot 37 \cdot 67 \quad (1.1)$$

and that it contains a unique conjugacy class of subgroups $\cong G_2(5)$. Let A_0 be one of them and B a Borel subgroup of A_0 , i.e. a 5-Sylow normalizer.

Then B is also a Borel group in A .

Furthermore, let $T \cong 4^2$ be a (maximal) torus in B , N_0 and N its normalizers in A_0 and A , respectively, and $W_0 = N_0/T$ and $W = N/T$ the corresponding Weyl groups.

From the theory of Chevalley groups, cf. e.g. Carter (1972), we deduce

$$W_0 \cong D_{12} \cong S_3 \times S_2 \tag{1.2}$$

while Kantor [1981] shows

$$W \cong S_4 \times S_3 . \tag{1.3}$$

A proper subgroup of A or A_0 which contains a Borel group will be called parabolic.

Kantor [1981] has shown that A contains exactly three conjugacy classes of maximal parabolic groups. They can be associated with the points P , lines L and planes F of a Tits geometry with the Buekenhout diagram

$$\begin{array}{ccccc}
 \circ & \text{-----} & \circ & \text{-----} & \overset{6}{\circ} & \text{-----} & \circ \\
 \delta & & 5 & & & & 5 \\
 P & & L & & & & F \\
 G_2(5) & & 5^{1+4} : 4S_6 & & & & 5^3 \cdot SL_3(5)
 \end{array} \tag{1.4}$$

Two objects (points, lines or planes) are called incident with each other if their intersection (as groups) is parabolic.

The apartment $A(T)$ associated with T is the set of all objects fixed by T . $A(T)$ is a subgeometry with Buekenhout diagram

$$\begin{array}{ccccc}
 \circ & \text{-----} & \circ & \text{-----} & \overset{6}{\circ} & \text{-----} & \circ \\
 1 & & 1 & & & & 1
 \end{array} \tag{1.5}$$

and may be represented (Kantor (1981)) by a simplicial complex A of dimension 2:




Fig. 1

Here the 0-, 1-, 2-simplices correspond to the 12 points, 36 lines, 24 planes of the apartment, respectively.

The Weyl group $W \cong S_4 \times S_3$ acts as S_4 on the numbers $\{1, 2, 3, 4\}$ and as S_3 on the letters $\{a, b, c\}$.

In analogy to Chevalley theory we now define the root groups associated with T as the groups X obeying the conditions:

- (1) $X \cong (\mathbb{F}_5, +) \cong 5$;
- (2) $T \leq N(X)$.

It follows from the known structure of $C_{Ly}(5B)$ (Lyons (1972)) that all root groups must be generated by $5A$ -elements.

Since $C_{Ly}(5A) \cong 5^{1+4} : (2 \wedge A_6)$ does not contain a Klein four group, $C_T(X) \cong 4$.

The A -normalizer of a $5A$ -group is a line. Thus there is a natural bijection between the root groups and the lines in $A(T)$.

The extension of T with the commutator subgroup

$$W' \cong A_4 \times A_3 \tag{1.6}$$

of W splits, so there is a unique element k of order 3 in N which corresponds to the Weyl element (abc) and centralizes T . In fact (Lyons [1972]),

$$C_A(T) = T \times K \tag{1.7}$$

with

$$K = \langle k \rangle \tag{1.8}$$

Furthermore, there is a set of 16 complements of T in the unique $T : A_4$ fixing the letters a, b, c . These groups are evidently conjugate under T , so we may elect an arbitrary one of them and denote it by Ω .

Ω is generated by 4 elements $\omega_i (1 \leq i \leq 4)$ which correspond to the 120° -rotations with centres in the points whose names contain the number i .

Each ω_i is uniquely determined by the choice of Ω and the corresponding Weyl permutation, namely

$$\omega_1 \rightarrow (234); \quad \omega_2 \rightarrow (143); \quad \omega_3 \rightarrow (124); \quad \omega_4 \rightarrow (132) \quad \text{in } W. \tag{1.9}$$

The group

$$\langle \omega_i, k \rangle = \Omega \times K \cong A_4 \times A_3 \tag{1.10}$$

(one of just 16 complements of T in $N' = T : W' \cong 4^2 : (A_4 \times A_3)$) is represented as a regular permutation group on the root groups X_L .

This allows us to specify a set of 36 generators (“roots”) for each of the 36 X_L .

We are free to take any generator for one of them, e.g. $X(1a, 2b)$. Call it $x(1a, 2b)$. Then apply $\Omega \times K$ to this root to define the remaining ones. A complete system of 36 root elements generated in this way will be called a standard (root) system.

Without restriction of generality we may assume that the Chevalley relations of $G_2(5)$ hold in the form described below (the exact exponents depend on the choice of $x(1a, 2b)$, but this clearly does not matter, because all allowed possibilities are equivalent and lead to the same group):

It will be convenient to define an orientation of the lines in \mathcal{A} according to the rules

$$a \rightarrow b, \quad b \rightarrow c, \quad c \rightarrow a. \tag{1.11}$$

Now we consider a point P in \mathcal{A} .

The 6 lines incident with P form a complete set of long roots for the stabilizer $\Lambda(P)$ of P , isomorphic to the Chevalley group $G_2(5)$, while the short roots are given by the sides of the (small) hexagon with centre P spanned by the long roots.

We denote the long and short roots by L_i and K_i ($i \in \mathbb{F}_7^*$), respectively, in the following manner:




Fig. 2

where L_i points from P to P_i for $i=1, 2, 4$ (squares in \mathbb{F}_7) and from P_i to P for $i=3, 5, 6$ (non-squares).

The 12 roots in Fig. 2 follow each other in the same order as they do in the standard G_2 root system.

Then the nontrivial Chevalley relations are

$$[L_i, L_{2i}] = L_{3i}^4, \tag{1.12}$$

$$[K_i, K_{3i}] = L_{2i}^3, \tag{1.13}$$

$$[K_i, K_{2i}] = L_{2i}^3 K_{3i}^3 L_{6i}^2, \tag{1.14}$$

$$[K_i, L_{4i}] = L_{2i}^4 K_{3i} L_{6i}^4 K_{2i}, \tag{1.15}$$

$$[L_i, K_{3i}] = K_{5i} L_{3i} K_i^4 L_{2i}^4 \tag{1.16}$$

combined with the information that for all $i \in \{1, 2, 4\}$ the mappings

$$K_i \Rightarrow \begin{vmatrix} 1 & 1 \\ \cdot & 1 \end{vmatrix}, \quad K_{-i} \Rightarrow \begin{vmatrix} 1 & \cdot \\ 4 & 1 \end{vmatrix}, \tag{1.17}$$

and

$$L_i \Rightarrow \begin{vmatrix} 1 & 1 \\ \cdot & 1 \end{vmatrix}, \quad L_{-i} \Rightarrow \begin{vmatrix} 1 & \cdot \\ 3 & 1 \end{vmatrix} \tag{1.18}$$

are isomorphisms from $\langle K_i, K_{-i} \rangle$ and $\langle L_i, L_{-i} \rangle$ onto $SL_2(5)$.

It should be noted that our relations differ slightly from those described, e.g., in Humphreys [1975]. This is due to our more symmetric choice of the roots which is more convenient in the context of the Lyons group.

For later reference, we construct an explicit $G_2(5)$ root system in the 7-dimensional minimal representation over \mathbb{F}_5 as the full automorphism group of the “septime” algebra with generators $e_i (i \in \mathbb{F}_7)$ and the skew-symmetric product given by

$$e_{i+j} \cdot e_{i+2j} = e_{i+4j} \quad (i \in \mathbb{F}_7; j \in \{1, 2, 4\}) . \tag{1.19}$$

Up to conjugacy in $G_2(5)$, our matrices are uniquely determined:

$$L_1 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & . & . & . & . & . \\ . & . & 1 & . & 3 & 1 & . \\ . & . & . & 1 & . & . & . \\ . & . & . & . & 1 & . & . \\ . & . & 2 & . & 1 & . & 1 \\ . & . & . & 4 & . & 1 & 2 \\ . & . & . & . & . & 4 & 3 & 1 \end{vmatrix}, \quad L_2 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & . & . & 2 & 1 & . \\ . & . & 1 & . & . & . & . \\ . & . & . & 1 & 4 & 2 & . \\ . & . & 3 & . & 1 & 1 & . \\ . & . & . & 4 & . & 3 & . & 1 \\ . & . & . & . & . & . & . & 1 \end{vmatrix}, \quad L_4 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & 3 & . & . & . & 1 \\ . & . & 2 & 1 & 1 & . & . \\ . & . & . & 4 & 1 & . & 3 \\ . & . & . & . & . & 1 & . \\ . & . & . & . & . & . & 1 \\ . & . & . & . & . & . & 1 \\ . & . & 4 & . & 2 & . & 1 \end{vmatrix}, \tag{1.20}$$

$$L_6 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & . & . & . & . & . \\ . & . & 1 & . & 1 & 3 & . \\ . & . & . & 1 & . & . & . \\ . & . & . & . & 1 & . & . \\ . & . & 4 & . & 1 & . & 3 \\ . & . & . & 2 & . & 1 & 4 \\ . & . & . & . & . & 2 & 1 & 1 \end{vmatrix}, \quad L_5 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & . & . & 4 & 3 & . \\ . & . & 1 & . & . & . & . \\ . & . & . & 1 & 2 & 4 & . \\ . & . & 1 & . & 3 & 1 & . \\ . & . & . & 2 & . & 1 & . & 1 \\ . & . & . & . & . & . & . & 1 \end{vmatrix}, \quad L_3 = \begin{vmatrix} 1 & . & . & . & . & . & . \\ . & 1 & 1 & . & . & . & 3 \\ . & . & 4 & 1 & 3 & . & . \\ . & . & . & 2 & 1 & . & 1 \\ . & . & . & . & . & 1 & . \\ . & . & . & . & . & . & 1 \\ . & . & . & . & . & . & 1 \\ . & . & 2 & . & 4 & . & 1 \end{vmatrix}, \tag{1.21}$$

$$K_1 = \begin{vmatrix} 1 & 1 & . & 2 & . & . & . \\ 4 & 3 & . & 4 & . & . & . \\ . & . & 1 & . & 1 & 3 & . \\ 3 & 4 & . & 4 & . & . & . \\ . & . & . & 4 & . & 1 & 2 \\ . & . & . & 2 & . & 1 & 1 \\ . & . & . & . & . & 3 & 4 & 1 \end{vmatrix}, \quad K_2 = \begin{vmatrix} 1 & . & 1 & . & . & . & 2 \\ . & 1 & . & . & 4 & 2 & . \\ 4 & . & 3 & . & . & . & 4 \\ . & . & . & 1 & 2 & 1 & . \\ . & . & 1 & . & 3 & 1 & . \\ . & . & 3 & . & 4 & . & 1 \\ 3 & . & 4 & . & . & . & 4 \end{vmatrix}, \quad K_4 = \begin{vmatrix} 1 & . & . & . & 1 & 2 & . \\ . & 1 & 1 & . & . & . & 3 \\ . & . & 4 & 1 & 2 & . & . \\ . & . & . & 3 & 1 & . & 4 \\ 4 & . & . & . & 3 & 4 & . \\ 3 & . & . & . & 4 & 4 & . \\ . & . & 2 & . & 1 & . & 1 \end{vmatrix}, \tag{1.22}$$

$$K_6 = \begin{vmatrix} 1 & 1 & . & 3 & . & . & . \\ 4 & 3 & . & 1 & . & . & . \\ . & . & 1 & . & 4 & 3 & . \\ 2 & 1 & . & 4 & . & . & . \\ . & . & 1 & . & 1 & . & 2 \\ . & . & . & 2 & . & 1 & 4 \\ . & . & . & . & . & 3 & 1 & 1 \end{vmatrix}, \quad K_5 = \begin{vmatrix} 1 & . & 1 & . & . & . & 3 \\ . & 1 & . & . & 1 & 2 & . \\ 4 & . & 3 & . & . & . & 1 \\ . & . & . & 1 & 2 & 4 & . \\ . & . & 4 & . & 3 & 1 & . \\ . & . & 3 & . & 1 & . & 1 \\ 2 & . & 1 & . & . & . & 4 \end{vmatrix}, \quad K_3 = \begin{vmatrix} 1 & . & . & . & 1 & 3 & . \\ . & 1 & 4 & . & . & . & 3 \\ . & . & 1 & 1 & 2 & . & . \\ . & . & . & 3 & 1 & . & 1 \\ 4 & . & . & . & 3 & 1 & . \\ 2 & . & . & . & 1 & 4 & . \\ . & . & 2 & . & 4 & . & 1 \end{vmatrix}. \tag{1.23}$$

Conjugation with $\Omega \times K$ merely permutes the roots (without exponents). Because of this fact, all standard systems are equivalent and lead to the same set of relations.

The lines in $A(T)$ form 3 parallel classes of 12 lines each. Every parallel class splits into 2 connected components called (great) circles.

A special line pair is a pair (L, L') of lines which are contained in a great circle and either have one point in common ("long" pair) or are mutual antipodes ("short" pair). The reason for this notation is that short and long special pairs form opposite pairs of short and long roots, respectively, in a certain $G_2(5)$ subgroup.

Since two opposite long root groups in $G_2(5)$ have the same centralizer in T , this must also be true for all 6 lines in a great circle.

The centralizer in the point $P \in A(T)$ of an arbitrary T -involution z contains 4 roots in each parallel class. Thus the group $\langle \Pi \rangle$ generated by a parallel class Π is a subgroup of $H = C_A(z) \cong 2 \wedge A_{11}$.

Let $\bar{H} = H/\langle z \rangle \cong A_{11}$.

As all subgroups isomorphic to 4^2 are conjugate in H we may assume without restriction that

$$\bar{T} = T/\langle z \rangle = \langle (1234)(5678), (1234)(8765) \rangle . \tag{1.24}$$

In H or \bar{H} exactly 12 groups $\cong 5$ exist which are normalized by T or \bar{T} . Except for a permutation of the letters $\{1, 2, 3, 4, 5, 6, 7, 8, 9, X, E\}$ normalizing T only the following correspondence between the roots and the permutations in \bar{H} is allowed by the Chevalley relations for the points:




Fig. 3

It is obvious that these roots generate $2 \wedge A_{11}$.

2. Definition and Simple Geometric Properties of the Group Γ

The results of Sect. 1 lead us to the definition of a group Γ as follows:

Γ is generated by 36 elements x_L , bijectively associated with the lines L in A (Fig. 1). The defining relations of Γ are

- (1) $C(P)$ -relations for every point P in A ;
- (2) $S(\Pi)$ -relations for each parallel class Π in A .

Here $C(P)$ is the set of Chevalley relations of $G_2(5)$ (cf. Sect. 1) while $S(\Pi)$ is an arbitrary system of defining relations for $2 \wedge A_{11}$. Best suited for our purpose are the Schur relations:

$$t_i^3 = 1 \quad (1 \leq i \leq 9) , \tag{2.1}$$

$$(t_i \cdot t_j)^2 = z \quad (1 \leq i, j \leq 9; i \neq j) , \tag{2.2}$$

$$z^2 = 1 \tag{2.3}$$

where the generator t_i corresponds to the permutation (iXE) .

We are now able to translate between the two sets of generators of $2 \wedge A_{11}$ (here $x(P, Q)$ is the root element which belongs to the line connecting the points P and Q):

$$t_9 = x(3c, 4a) \cdot x(4b, 3c)^{-1} \cdot x(3a, 4b) \cdot x(4a, 3b)^{-1} \cdot x(3a, 4b) \cdot x(4b, 3c) \cdot x(3c, 4a)^{-1} , \tag{2.4}$$

$$t_1 = x(3a, 4b)^{-1} \cdot t_9 \cdot x(3a, 4b) , \quad t_2 = x(3a, 4b)^2 \cdot t_9 \cdot x(3a, 4b)^{-2} , \tag{2.5}$$

$$t_3 = x(3a, 4b) \cdot t_9 \cdot x(3a, 4b)^{-1} , \quad t_4 = x(3a, 4b)^{-2} \cdot t_9 \cdot x(3a, 4b)^2 , \tag{2.6}$$

$$t_5 = x(1a, 2b)^{-1} \cdot t_9 \cdot x(1a, 2b) , \quad t_6 = x(1a, 2b)^2 \cdot t_9 \cdot x(1a, 2b)^{-2} , \tag{2.7}$$

$$t_7 = x(1a, 2b) \cdot t_9 \cdot x(1a, 2b)^{-1} , \quad t_8 = x(1a, 2b)^{-2} \cdot t_9 \cdot x(1a, 2b)^2 . \tag{2.8}$$

The reverse transformations are:

$$x(3a, 4b) = t_9^{-1} \cdot t_1^{-1} \cdot t_4 \cdot t_2^{-1} \cdot t_3 \cdot t_1 \cdot t_9 , \quad x(1a, 2b) = t_9^{-1} \cdot t_5^{-1} \cdot t_8 \cdot t_6^{-1} \cdot t_7 \cdot t_5 \cdot t_9 , \tag{2.9}$$

$$x(4b, 3c) = t_2 \cdot t_4 \cdot t_1^{-1} \cdot t_3 \cdot t_2^{-1} , \quad x(2b, 1c) = t_6 \cdot t_8 \cdot t_5^{-1} \cdot t_7 \cdot t_6^{-1} , \tag{2.10}$$

$$x(3c, 4a) = t_1^{-1} \cdot t_4 \cdot t_2^{-1} \cdot t_3 \cdot t_1 , \quad x(1c, 2a) = t_5^{-1} \cdot t_8 \cdot t_6^{-1} \cdot t_7 \cdot t_5 , \tag{2.11}$$

$$x(4a, 3b) = t_9^{-1} \cdot t_4^{-1} \cdot t_1 \cdot t_3^{-1} \cdot t_2 \cdot t_4 \cdot t_9 , \quad x(2a, 1b) = t_9^{-1} \cdot t_8^{-1} \cdot t_5 \cdot t_7^{-1} \cdot t_6 \cdot t_8 \cdot t_9 , \tag{2.12}$$

$$x(3b, 4c) = t_3 \cdot t_1 \cdot t_4^{-1} \cdot t_2 \cdot t_3^{-1} , \quad x(1b, 2c) = t_7 \cdot t_5 \cdot t_8^{-1} \cdot t_6 \cdot t_7^{-1} , \tag{2.13}$$

$$x(4c, 3a) = t_4^{-1} \cdot t_1 \cdot t_3^{-1} \cdot t_2 \cdot t_4 , \quad x(2c, 1a) = t_8^{-1} \cdot t_5 \cdot t_7^{-1} \cdot t_6 \cdot t_8 . \tag{2.14}$$

By the main result of Meyer et al. [1985] the Lyons group possesses a 111-dimensional irreducible representation over \mathbb{F}_5 . In this we can easily identify 36 elements which generate Ly and satisfy all relations defining Γ .

Hence we have

- Lemma 1.** (a) *The Lyons group is a homomorphic image of Γ ;*
 (b) *Γ has a 111-dimensional nontrivial representation over \mathbb{F}_5 .*

Let us now consider the following subconfigurations of A in Fig. 1:




Fig. 4a-c






The subgroups of Γ generated by the lines in Fig. 4a-c (respectively) are called $\Gamma(P)$; $\Gamma(L)$; $\Gamma(F)$.

For any set $\{\mathcal{O}, \mathcal{O}', \dots\}$ of objects we define $\Gamma(\mathcal{O}, \mathcal{O}', \dots)$ as the intersection of the groups $\Gamma(\mathcal{O}), \Gamma(\mathcal{O}'), \dots$

With the above notation we have

- Theorem 1.** (a) $\Gamma(P) \cong G_2(5)$;
 (b) $\Gamma(L) \cong 5^{1+4} : (2 \wedge A_6)$;
 (c) $\Gamma(F) \cong 5^3 \cdot SL_3(5)$.

Theorem 2. *The lines in the following configurations (omitting the dotted lines):*



- (f) great circle
 (g) parallel class

generate groups which are isomorphic to:

- (a) 5 ;
- (b) 5^3 ;
- (c) 5^{1+4} ;
- (d) $2 \wedge A_6$;
- (e) $5 \times (2 \wedge A_6)$;
- (f) $2 \wedge A_7$;
- (g) $2 \wedge A_{11}$.

Proof of Theorems 1 and 2. (1a) Due to the $C(P)$ -relations, $\Gamma(P)$ is a homomorphic image of $G_2(5)$ (using a theorem of Steinberg, cf. Carter (1972), Theorem 12.1.1), so $\Gamma(P)$ is $\cong G_2(5)$ or $\cong 1$.

In the latter case a root group in $\Gamma(P)$ and hence also in $\Gamma(P')$ for a neighbouring point P' of P would be trivial, so $\Gamma(P') = 1$, too. This leads to $\Gamma = 1$, contradicting Lemma 1.

(2g) Because of the $S(\Pi)$ -relations, $\langle \Pi \rangle$ is a homomorphic image of $2 \wedge A_{11}$, so is $\cong 2 \wedge A_{11}, A_{11}$ or 1. Only the first possibility is in conformity with (1a), since a long special line pair generates $SL_2(5) \cong 2 \wedge A_5$. The remaining statements in Theorem 2 now follow immediately from (1a) and (2g).

(1b) Since (2c) and (2d) hold, we need only show that 5^{1+4} is normalized by $2 \wedge A_6$. This follows from (1a), applied to the two points incident with L .

(1c) Analogously to (1b), we conclude with the help of (2b) that the three lines incident with F generate a normal subgroup $\Gamma_0(F) \cong 5^3$ of $\Gamma(F)$.

The images in $\Gamma(F)/\Gamma_0(F)$ of the root subgroups in $\Gamma(F)$ fulfill all of the Chevalley relations for the group $SL_3(5)$ (which is defined by these relations) if we map them as follows:




Fig. 5

$SL_3(5)$ is simple, and according to (2c) not all of the images can be trivial; thus $\Gamma(F) \cong 5^3 \cdot SL_3(5)$. This extension does not split, since $\Gamma(F)$ contains a 5-Sylow subgroup of $G_2(5)$ and therefore elements of order 25. This establishes Theorems 1 and 2.

We now define for an arbitrary (long or short) special line pair L, L' the groups $T_{LL'}$ and $Q_{LL'}$ as follows:

Let $T_{LL'}$ be the common normalizer of the root groups L and L' and let $Q_{LL'}$ be the set normalizer of $\{L, L'\}$ in $\langle L, L' \rangle \cong SL_2(5)$.

Furthermore, for each great circle K and each parallel system Π we introduce the abbreviations

$$T_K = \langle T_{LL'} : L, L' \text{ special line pair in } K \rangle, \tag{2.15}$$

$$Q_K = \langle Q_{LL'} : L, L' \text{ special line pair in } K \rangle, \tag{2.16}$$

$$T_\Pi = \langle T_{LL'} : L, L' \text{ special line pair in } \Pi \rangle, \tag{2.17}$$

$$Q_\Pi = \langle Q_{LL'} : L, L' \text{ special line pair in } \Pi \rangle, \tag{2.18}$$

as well as

$$T = \langle T_{LL'} : L, L' \text{ special line pair} \rangle, \tag{2.19}$$

$$Q = \langle Q_{LL'} : L, L' \text{ special line pair} \rangle \tag{2.20}$$

and for any point P :

$$T(P) = \langle T_{LL'} : L, L' \text{ special line pair in } \Gamma(P) \rangle. \tag{2.21}$$

Of course, $T(P)$ is the torus in $\Gamma(P) \cong G_2(5)$ belonging to our apartment. We next show

- Theorem 3.** (a) For a special line pair L, L' in the circle K we have $T_{LL'} = T_K \cong 4$;
 (b) for every parallel class Π : $T_\Pi \cong 4 \times 2$;
 (c) for all points P : $T(P) = T \cong 4^2$.

Proof. (a) and (b) follow from an easy calculation in $\langle \Pi \rangle \cong 2 \wedge A_{11}$. Trivially, we have $T(P) \leq T$. With (a) we deduce for every great circle K with an arbitrary but fixed P that $T_K \leq T(P)$. Since $\langle T_K \rangle = T$, we get (c).

- Theorem 4.** (a) For all special line pairs L, L' : $Q_{LL'} = N_{\langle L, L' \rangle}(T_{LL'}) \cong Q_8$, the quaternion group of order 8; the intersection of T with $Q_{LL'}$ is $T_{LL'}$;
 (b) T is a normal subgroup of Q ;
 (c) each element q of Q permutes the lines of A , inducing an automorphism of A as a simplicial complex;
 (d) the image of this action is the full automorphism group $S_4 \times S_3$ of A .

Proof. The first part of (a) is immediate since $\langle L, L' \rangle \cong SL_2(5)$. The second part can be verified in $\Gamma(P)$ for an appropriate point P .

In this $\Gamma(P)$ we also see that $Q_{LL'}$ normalizes $T(P) = T$, thus the same holds true for $Q = \langle Q_{LL'} \rangle$. Furthermore, each $T_{LL'}$ is contained in $Q_{LL'}$, hence in Q ; so $T = \langle T_{LL'} \rangle$ is a subgroup of Q . This proves (b).

Let q be an element of $Q_{LL'}$. If q is contained in $T_{LL'} < T$, (c) holds trivially. If q is in $Q_{LL'} \setminus T_{LL'}$, q induces a permutation of the groups of order 5 which are normalized by T in each of the groups $\Gamma(P)$ and $\langle \Pi \rangle$ where P is any point with $L, L' < \Gamma(P)$ and Π the parallel system containing L and L' . But all these groups of order 5 are root groups.

From the $C(P')$ -relations for appropriate points P' we find that the 16 remaining roots are also permuted. Inspection of the permutations generated by Q easily leads to (c) and (d).

3. Some Geometric Subgroups of Γ

Let Π be a parallel class and P a point in A . The group $H = \langle \Pi \rangle$ is $\cong 2 \wedge A_{11}$ by Theorem 2.g. We denote the unique involution in $Z(H)$ by z .

We now prove

Theorem 5. *The intersection of H and $\Gamma(P)$ is $C_{\Gamma(P)}(z) \cong (1/2) \cdot 2 \wedge (S_5 \times S_5)$.*

Proof. Since all pairs (Π, P) are equivalent under Q (Theorem 4.d), we may restrict ourselves to the case $P = 1a$ and $\Pi =$ parallel system of Fig. 3. Then H and $\Gamma(P)$ obviously contain the 4 roots $x(1a, 2b), x(2c, 1a), x(3b, 4c), x(4b, 3c)$ which generate a group $SL_2(5) \vee SL_2(5) \cong 2 \wedge (A_5 \times A_5)$ of index 2 in $C_{\Gamma(P)}(z) \cong (1/2) \cdot 2 \wedge (S_5 \times S_5)$. This group is enlarged by $T - T < H$ and $T < \Gamma(P)$ because of Theorem 3.c – to the full centralizer of z in $\Gamma(P)$. As z is in the centre of H , the intersection of H and $\Gamma(P)$ is a subgroup of $C_{\Gamma(P)}(z)$; hence the proposition.

We want to consider several groups which are defined symmetrically with respect to the apartment $A(T)$.

Let

$$U_1 = \Gamma(1a, 1b, 1c) \ , \quad U_2 = \Gamma(2a, 2b, 2c) \ , \quad (3.1)$$

$$U_3 = \Gamma(3a, 3b, 3c) \ , \quad U_4 = \Gamma(4a, 4b, 4c) \ , \quad (3.2)$$

and

$$U = \langle U_1, U_2, U_3, U_4 \rangle \ . \quad (3.3)$$

It will be convenient to have a systematic notation for the circles, parallel systems and corresponding $2 \wedge A_{11}$ -subgroups in Γ :

We denote the circle containing the points with numbers i and j by K_{ij} and the parallel system consisting of K_{ij} and K_{kl} by $\Pi_{ij,kl}$. The corresponding T -involution will be called $z_{ij,kl}$, and we set $H_{ij,kl} = \langle \Pi_{ij,kl} \rangle$.

Hence the torus elements $z_{12,34}, z_{13,24}, z_{14,23}$ are canonically associated with the double transpositions in the symmetric group S_4 , while the circles $K_{12}, K_{13}, K_{14}, K_{23}, K_{24}, K_{34}$ belong to the transpositions of S_4 .

Let us now investigate the groups $U_i (1 \leq i \leq 4)$ and U :

Theorem 6. (a) $U_1 \cong U_2 \cong U_3 \cong U_4 \cong U_3(3)$;

(b) $U' = U; U/Z(U) \cong U_4(3) \cong O_6^-(3); Z(U) \leq 4 \times 3^2$.

Proof. We define

$$a = x(4b, 3c)^4 x(3b, 4c)^2 \Rightarrow (132) \text{ in } H_{12,34} \ , \quad (3.4)$$

$$b = x(4b, 3c)^1 x(3b, 4c)^3 \Rightarrow (143) \text{ in } H_{12,34} \ , \quad (3.5)$$

$$c = x(4b, 2c)^1 x(2b, 4c)^3 \Rightarrow (124) \text{ in } H_{13,24} \ , \quad (3.6)$$

$$r = x(1a, 2b)^1 x(2a, 1b)^3 \Rightarrow (568) \text{ in } H_{12,34} \ , \quad (3.7)$$

$\Gamma(1a), \Gamma(1b), \Gamma(1c)$ contain the $2 \wedge A_5$ -groups $\langle x(3b, 4c), x(4b, 3c) \rangle, \langle x(3c, 4a), x(4c, 3a) \rangle, \langle x(3a, 4b), x(4a, 3b) \rangle$ of $H_{12,34}$, acting on the sets $\{1, 2, 3, 4, X\}, \{1, 2, 3, 4, E\}, \{1, 2, 3, 4, 9\}$, respectively.

Their intersection, the $2 \wedge A_4$ -group on $\{1, 2, 3, 4\}$, is thus contained in $\Gamma(1a, 1b, 1c) = U_1$.

Obviously, analogous results for $H_{13.24}$ and $H_{14.23}$ hold. Hence, by (3.4), (3.5), (3.6),

$$\langle a, b, c \rangle \leq U_1 = \Gamma(1a, 1b, 1c) \leq \Gamma(1a) . \tag{3.8}$$

In $\Gamma(1a)$ we easily verify – see (1.20), ..., (1.23) – that

$$a^3 = b^3 = c^3 = 1 , \tag{3.9}$$

$$aba = bab , \quad aca = cac , \quad bcb = cbc , \tag{3.10}$$

$$a^b c^{-1} a^b = c^{-1} a^b c^{-1} ; \quad b^a c^{-1} b^a = c^{-1} b^a c^{-1} . \tag{3.11}$$

These relations form a presentation of the finite simple group $U_3(3)$, cf. Aschbacher and Hall [1973].

Since $\langle a, b, c \rangle$ is nontrivial, we deduce

$$U_3(3) \cong \langle a, b, c \rangle \leq U_1 \leq \Gamma(1a) \cong G_2(5) . \tag{3.12}$$

By inspection of the maximal subgroups of $G_2(5)$ we are left with three candidates for U_1 , namely $\langle a, b, c \rangle \cong U_3(3)$, $N_{\Gamma(1a)}(\langle a, b, c \rangle) \cong G_2(2)$ and $\Gamma(1a) \cong G_2(5)$ (Conway et al. [1985]).

But, by Theorem 5, $C_{U_1}(z_{12.34})$ is the intersection of U_1 and $H_{12.34}$, hence $C_{U_1}(z_{12.34}) = \langle a, b, T \rangle \cong 4S_4$.

$G_2(2)$ and $G_2(5)$ do not contain involution centralizers of this form, so

$$U_1 = \langle a, b, c \rangle \cong U_3(3) . \tag{3.13}$$

Since U_1, U_2, U_3, U_4 are conjugate to each other under Q , (a) follows. Furthermore, r and c are both contained in $\Gamma(3a)$ where we immediately establish the relations

$$r^3 = 1 , \quad rcr = crc \tag{3.14}$$

while in $H_{12.34} \cong 2 \wedge A_{11}$ the elements a and b evidently commute with r :

$$ra = ar , \quad rb = br . \tag{3.15}$$

By a result of Aschbacher and Hall [1973] the relations (3.9, 3.10, 3.11, 3.14, 3.15) form a presentation of the full Schur cover of the finite simple group $U_4(3) \cong O_6^-(3)$, so with the abbreviation

$$U_0 = \langle a, b, c, r \rangle \tag{3.16}$$

we get (because $U_4(3)$ is simple and $U_0 \neq 1$)

$$U'_0 = U_0 , \quad U_0/Z(U_0) \cong U_4(3) \tag{3.17}$$

and $Z(U_0)$ is a factor of the Schur multiplier $12 \times 3 \cong 4 \times 3^2$ of $U_4(3)$. To complete the proof of our theorem it remains to show that $U_0 = U$. First we have $a, b, c \in U_1 \leq U$ and $r \in U_3 \leq U$, so $U_0 \leq U$.

The reverse inequality amounts to $U_i \leq U_0$ for all $i \in \{1, 2, 3, 4\}$. Clearly this is true for $U_1 = \langle a, b, c \rangle$.

The intersection of U_1 and U_3 contains the torus T as well as c . Since $\langle c, T \rangle \cong 4S_4$ is maximal in U_3 and centralizes $z_{13.24}$, while $r \in U_3$ does not, we get

$$U_3 = \langle c, T, r \rangle \leq U_0 . \tag{3.18}$$

Let now $i=2$ or 4 . The intersections of U_i with U_1 and U_3 are different maximal subgroups ($\cong 4S_4$) of $U_i \cong U_3(3)$ and therefore they together generate U_i . Since they are contained in $\langle U_1, U_3 \rangle \leq U_0$, this completes the proof of the required equality

$$U = \langle a, b, c, r \rangle \tag{3.19}$$

at the same time establishing the theorem.

Having chosen a suitable unitary basis, the matrices in $SU_4(3)$ corresponding to the elements a, b, c, r are found to be

$$a = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1-i & -1+i \\ 0 & 0 & 1+i & 1+i \end{pmatrix}, \quad b = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1-i & 1-i \\ 0 & 0 & -1-i & 1+i \end{pmatrix}. \tag{3.20}$$

$$c = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1-i & 0 & -1+i \\ 0 & 0 & 1 & 0 \\ 0 & 1+i & 0 & 1+i \end{pmatrix}, \quad r = \begin{pmatrix} 1+i & -1-i & 0 & 0 \\ 1-i & 1-i & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{3.21}$$

where i is a square root of -1 in \mathbb{F}_9 .

The matrices in $SO_6^-(3)$ are given by

$$a = \begin{pmatrix} 1 & . & . & . & . \\ . & 1 & . & . & . \\ . & . & 1 & 1 & 2 & 2 \\ . & . & 2 & 1 & 1 & 2 \\ . & . & 1 & 2 & 1 & 2 \\ . & . & 1 & 1 & 1 & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 1 & . & . & . & . \\ . & 1 & . & . & . \\ . & . & 1 & 1 & 1 & 1 \\ . & . & 2 & 1 & 2 & 1 \\ . & . & 2 & 1 & 1 & 2 \\ . & . & 2 & 2 & 1 & 1 \end{pmatrix}. \tag{3.22}$$

$$c = \begin{pmatrix} 1 & 2 & . & . & 1 & 1 \\ 1 & 1 & . & . & 1 & 2 \\ . & . & 1 & . & . & . \\ . & . & . & 1 & . & . \\ 2 & 2 & . & . & 1 & 2 \\ 2 & 1 & . & . & 1 & 1 \end{pmatrix}, \quad r = \begin{pmatrix} 1 & . & . & . & . \\ . & 1 & . & . & . \\ . & . & 1 & 2 & 2 & 2 \\ . & . & 1 & 1 & 2 & 1 \\ . & . & 1 & 1 & 1 & 2 \\ . & . & 1 & 2 & 1 & 1 \end{pmatrix}. \tag{3.23}$$

4. The Lyons Group as a Homomorphic Image of Γ

We now define elements a, b, c, d, x in Γ by

$$a = \tau \cdot x(3b, 4c)^4 \cdot x(4b, 3c)^2, \tag{4.1}$$

$$b = x(3b, 4c)^2, \tag{4.2}$$

$$c = x(4b, 2c)^3, \tag{4.3}$$

$$d = x(1a, 2b) \cdot x(2c, 1a)^3 \cdot x(1a, 2b), \tag{4.4}$$

$$x = x(3b, 4c) \cdot x(4b, 3c) \cdot x(3b, 4c) \cdot \tau' \cdot x(4a, 3b) \cdot x(3b, 4c)^3 \cdot x(4a, 3b) \tag{4.5}$$

where τ and τ' are the torus elements

$$\tau = x(4b, 2c) \cdot x(2b, 4c)^4 \cdot x(4b, 2c)^2 \cdot x(2b, 4c)^2, \tag{4.6}$$

$$\tau' = x(1a, 4b)^4 \cdot x(4c, 1a) \cdot x(1a, 4b)^2 \cdot x(4c, 1a)^2. \tag{4.7}$$

We verified by computer that the images $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{x}$ of a, b, c, d, x in the 111-dimensional representation (cf. Lemma 1) obey all of the relations of Sims [1973], and hence they generate the Lyons group.

Furthermore,

$$\langle \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} \rangle \cong G_2(5) \tag{4.8}$$

while

$$\langle a, b, c, d \rangle \leq \Gamma(1a) \cong G_2(5). \tag{4.9}$$

Therefore

$$\langle a, b, c, d \rangle = \Gamma(1a) \tag{4.10}$$

$x \in Q$ by Theorem 4.c permutes the 36 root groups and corresponds to the automorphism (12)(34) · (ac) of the apartment.

$\langle a, b, c, d, x \rangle$ contains the 12 root groups in $\Gamma(1a)$ and, e.g.,

$$X(1a, 2b)^x = X(1b, 2c). \tag{4.11}$$

Because of the Chevalley relations these 13 root groups generate Γ . This shows the validity of

- Theorem 7.** (a) $\langle a, b, c, d, x \rangle = \Gamma$;
 (b) L_y is a homomorphic image of Γ .

Remark. To prove a relation in any subgroup Δ of Γ which is isomorphic to its image $\tilde{\Delta}$ in the representation, it is sufficient to check this relation for the appropriate 111-dimensional \mathbb{F}_5 -matrices.

In particular this holds true for the Sims relations which are expressed in elements of Δ alone.

We may apply this to the following three subgroups:

$$\Delta_x = \Gamma(1a) \cong G_2(5), \tag{4.12}$$

$$\Delta_c = H_{12,34} \cong 2 \wedge A_{11}, \tag{4.13}$$

$$\Delta_d = \langle \Gamma(L(2c, 1a)), T \rangle \cong 5^{1+4} : 4S_6. \tag{4.14}$$

The isomorphisms $\Delta_x \cong \tilde{\Delta}_x$ and $\Delta_c \cong \tilde{\Delta}_c$ have been verified in Theorems 1.a and 2.c, respectively.

$\Delta_d \cong \tilde{\Delta}_d$ follows immediately from Theorem 2.e and the fact that all 36 root groups are normalized by T .

These arguments suffice to prove the validity of all Sims relations except three.

We believe that the remaining relations also follow from our presentation of Γ , but we have not yet been able to show this.

5. Summary

The goal of this paper is to construct a root system for the Lyons group L_y in analogy to those of the Chevalley groups.

We make ample use of geometric properties of Ly .

We are confident that similar ideas can be applied to other (all ?) sporadic groups as well, perhaps in the long run leading to an understanding of these peculiar structures.

Concerning the geometry of the Lyons group itself, more information may be gained by a careful study of the 111-dimensional minimal representation over \mathbb{F}_5 .

Some initial results in that direction have been obtained.

We hope to present them – together with a proof of the isomorphism of the group Γ (defined in Sect. 2) with Ly – in the near future.

Acknowledgements. We are deeply indebted to Prof. Dr. Joachim Neubüser (RWTH Aachen) and his collaborators who supplied us in a most generous way with informations on several finite groups, character tables etc.

Without their friendly support and encouragement this investigation could not have been carried through.

References

- Aschbacher, M., Hall, M.: Groups generated by a class of elements of order 3. *J. Alg.* **24**, 591–612 (1973)
- Carter, R.: *Simple groups of Lie type*. New York: Wiley Interscience 1972
- Conway, J.H., Curtis, R.T., Norton, S.P., Parker, R.A., Wilson, R.A.: *Atlas of finite groups*. Oxford: Clarendon Press 1985
- Humphreys, J.E.: *Linear algebraic groups*. (Graduate Texts in Mathematics, Vol. 21). Berlin Heidelberg New York: Springer 1975
- Kantor, W.: Some geometries that are almost buildings. *Eur. J. Comb.* **2**, 239–247 (1981)
- Lyons, R.: Evidence for a new finite simple group. *J. Alg.* **20**, 540–569 (1972)
- Meyer, W., Neutsch, W.: Über 5-Darstellungen der Lyonsgruppe. *Math. Ann.* **267**, 519–535 (1984)
- Meyer, W., Neutsch, W., Parker, R.: The minimal 5-representation of Lyons' sporadic group. *Math. Ann.* **272**, 29–39 (1985)
- Sims, C.C.: The existence and uniqueness of Lyons' group, in: *Finite Groups '72* (Gainesville Conference), 138–141, Amsterdam: North-Holland 1973
- Wilson, R.A.: The subgroup structure of the Lyons group. *Math. Proc. Cambr. Philos. Soc.* **95**, 403–409 (1984)
- Wilson, R.A.: The maximal subgroups of the Lyons group. *Math. Proc. Cambr. Philos. Soc.* **97**, 433–436 (1985)
- Woldar, A.J.: On the maximal subgroups of Lyons' group. *Comm. Alg.* **15**, 1195–1203 (1987)

Received April 1, 1988; in revised form June 13, 1988

