

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0020

LOG Titel: XIII. Flos prunae Damascenae

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

tantur, mense Julio fulva producitur inde phalæna, cujus alarum superior quælibet albâ maculâ distingritur, sicuti figura habet,

XI. *Cerasus austera, florens.*

Hujuscemodi flavo quodam lucido tinctas erucas sæpè sæpiùs his super arboribus inveni, folia, quibus involvuntur, complicant, pilosæ sunt admodum tam antrosùm quàm retrorsùm gradientes, tactæ filo ex ore protenso descendunt, in sextum usquè diem mensis Maii eas enutrivì, telam exinde albam conficiebant, & in nymphas fuscas immutatæ vigesimâ septimâ die Octobris, in papiliones è lucido fuscis, ad latus depictos, migrarunt. Tales in cerasis quàm plurimas deprehendi subflavas erucas, folia viridia complicant, ac pilosæ admodum sunt tam antrosùm quàm retrorsùm gradientes, tactæ filo ex ore protenso descendunt, quo mediante pariter reascendere noverunt, in sextum usque diem mensis Maii eas enutrivì, tunc albam telam conficiebant, vigesimâ septimâ die Octobris è lucido flavo tinctæ colore phalænulæ prodibant, qualis in folio viridi suprâ annotata est.

XII. *Viola Lutea.*

Super hocce murario vel potius montano flore hujusmodi tenues erucæ virides inveniuntur, aliis erucis proportionatè longiores, sub corpore pedibus carent, progredientes corporis partem posticam priori jungunt, gibbum hocce modo formantes, iter suum accelerant; animadverti, quod mense Junio folio sese firmiter intexentes in aureliam fuscam immutentur, Julio fuscus & elegans inde papilio prodit, qualis proximè florem hunc est appictus.

Flori flavo superjacet vermiculus, quem in rapâ reperi, hic in doliolum immutabatur, decem diebus elpasis parva quædam musca, in arbore delineata, prodiit.

XIII. *Flos prunæ Damascenæ.*

Formosam hanc virescentem striis nigris, coralliisq; flavis adornatam erucam hisce foliis in mensẽ usque Junium-

enutrivi, ubi argenti more resplendenti & instar chartæ pergamænæ Tabulosæ ovalis figuræ telæ sese intexuit, & in aureliam fuscâ defigurabatur, in principio mensis Augusti elegans inde phalæna prodiit, cujus alæ inferiores flavæ fusco colore striatæ, sicciterque maculatæ sunt.

Inferiori folio virenti, crustâ fuscâ eâque solidâ munitum adhæret animalculum, hoc domunculam suam cochlearum adinstar secum circumfert tardigradum, hisce foliis in vigesimum usque Junii diem illud enutrivi, exinde non movebatur, undecimo die Julii, qualis supra eruculam fuscâ repræsentatur, phalæna prodiit alba, aliud è regione animalculum ejusdem est naturæ.

XIV. *Grossularia hortensis non Spinosa florens.*

HAc in arbore genus aliquod erucarum animadvertitur, quarum anterior pars corporis flava, posterior autem superius alba & inferius flava est, eas, in aurelias usquedùm immutarentur, infantem fasciis involutum apprimè referentes, ac auro velut & argento coruscas, enutrivi, in fine Junii papilio prodiit, cujus latera superius summè flava, fusco colore maculata, & inferius fusca, maculis nigris adornata sunt.

XV. *Ranunculus pratensis.*

HUjuscemodi discolors erucas in mensem usque Junii hæc fronde nutrivi, tunc in aureliam nigram migrabant, undè papilio prodiit elegans, cujus alæ superiores è lucido flavæ punctis nigris aspersæ, sed inferiores auriaco colore, virgisque nigris adornatæ erant.

XVI. *Cærefolium.*

HÆc herba virentis, albæ, virgulatæ, glabræ erucæ pabulum est, in finem usque Maii enutriva tenuem sibi telam parat, quâ confectâ, in aureliam spadicei coloris immutatur, diebus quatuordecim elapsis fusca phalæna producta est, qualem figura repræsentat.

XVII.