

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0024
LOG Titel: XVII. Vermes miraculosi
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XVII. *Vermes miraculosi.*

DUas has ingentes, formâ tamen, atque colore disconvenientes erucas in herbâ, earundem pabulo reperi, simili cum aliis Metamorphose mutabiles, quæ semetipsas intexentes, in aurelias cineritii coloris immutantur, è quibus ingentes duæ obscuro quodam auriaco colore præditæ phalænæ postmodum, in totum sibi propemodum similes, in conspectum venerunt.

XVIII. *Malus mellea florens.*

HOC erucarum genus arbori cuilibet fructiferæ summè nocivum est, interdiù per universam arborem disperguntur, noctù verò in fasciculum aranearum telæ similem unâ conve-niunt, hæc eruca subfuscæ est, cuilibet infectioni dorsi cæruea, simul ac rubicunda corallia sunt intertexta, telâ tenui circumcircâ confectâ, in aureliam fuscam invertuntur, undè diebus quatuordecim elapsis albicans fuscis striis distincta phalæna prorumpit, ovula parta Lanugine subflavâ contegunt, ita ut in conspectum non veniant, hocce modo hyemali tempore à frigore sese tuentur. In pomis itidem subflavam eruciam inveni, quæ sub finem mensis Maii telæ sese intexit, in quâ hyemali tempore morabatur, verno, qualis suprà virentia folia conspicitur, exigua musca prorupit.

XIX. *Rosa incarnata.*

SUPER rosæ calyce viridem eruculam sedentem conspicies, quæ non folia solummodo viridia depascitur, sed & caly-cem excavat, ità ut rosarum accretionem impedit, in fine mensis Maii in Aureliam ochræ colore tintam immutatur, Junio papiliones in rosis appicti prodeunt.

Subtus rosæ caulem erucula fusca prorept, quæ medios caly-ces per modum prioris exedit, Majo metamorphosin subit, Junio mense, qualis ad latus volitans appictus est, papilio prodit, hujus alæ superiores ochræ, inferiores verò cinereum colorem referunt.

Infe-