

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0025

LOG Titel: XVIII. Malus mellea florens

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XVII. *Vermes miraculofi.*

DUas has ingentes, formâ tamen, atque colore disconvenientes erucas in herbâ, earundem pabulo reperi, fimilium cum aliis Metamorphose mutabiles, quæ femetiplus intextentes, in aurelias cineritii coloris immutantur, è quibus ingentes duæ obscuro quodam auriaco colore præditæ phalænæ postmodum, in totum sibi propemodùm similes, in conspectum venerunt.

XVIII. *Malus melleâ florens.*

HOC erucarum genus arbori cuilibet fructiferæ summè nocivum est, interdium per universam arborem disperguntur, noctû verò in fasciculum aranearum telæ similem unâ conveniunt, hæc eruca subfusca est, cuilibet infectioni dorfi cærulea, simul ac rubicunda corallia sunt intertexta, telâ tenui circumcircâ confectâ, in aureliam fuscâ invertuntur, undè diebus quatuordecim elapsis albicans fuscis striis distincta phalæna prorumpit, ovula parta Lanugine subflavâ contegunt, ita ut in conspectum non veniant, hocce modo hyemali tempore à frigore sese tuentur. In pomis itidem subflavam eruculam inveni, quæ sub finem mensis Maii telæ sese intexuit, in quâ hyemali tempore morabatur, verno, qualis suprâ virentia folia conspicitur, exigua musca prorupit.

XIX. *Rosa incarnata.*

SUPER rosæ calyce viridem eruculam sedentem conspicias, quæ non folia solummodò viridia depascitur, sed & calycem excavat, ita ut rosarum accretionem impediât, in fine mensis Maii in Aureliam ochræ colore tinctam immutatur, Junio papiliones in rosis appicti prodeunt.

Subtus rosæ caulem erucula fusca prorepat, quæ medios calyces per modum prioris exedit, Majo metamorphosin subit, Junio mense, qualis ad latus volitans appictus est, papilio prodit, hujus alæ superiores ochræ, inferiores verò cinereum colorem referunt.

Infe-