

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0029

LOG Titel: XXII. Rosa multiplex, media

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

Inferius folio viridi viridis eruca infidet, sub foliis supina pabulum fumere solita, in decimam usque Junii diem eam enutriti, ubi perfectâ tenui, albâque texturâ in aureliam lignei coloris immutabatur, vigesimâ sextâ ejusdem inde phalæna fusca prodiit, qualis in Marcescente rosâ denotatur.

XX. *Oryacantha florens.*

Diversicolor hæc eruca ut plurimum in dumis invenitur, neque tamen ideò à pomorum foliis abhorret, à me quodam tempore enutrita telam cineritii coloris sibi texuit, & in aureliam fuscâ immutabatur, quæ vivens tanti sensûs erat, ut tacta sese circumvolveret, diebus duodecim præterlapsis alba prodiit inde phalæna, quæ flava pariens ovula extincta est.

Sordidos hosce vermes in cloacis graveolentibus est reperire, mutantur in doliola caudati, undè etiam muscoli nuncupantur, spatio septendecim dierum elapso muscam educi sub flavam inertem animadverti.

XXI. *Cotoneæ flos.*

Tantas his in arboribus erucas inveni flavo ochræ colore tinctas, striis nigris, punctisque albicantibus asperfas, cauda, caputque auriaci sunt coloris, tactæ multâ motitatione caput agitant, hisce foliis eas enutriti, usquedùm sub flavam tabulosamque texturam conficientes in Aurelias immutarentur, trium hebdomadarum spatio ingens ochræ colore tincta phalæna prodiit quæ totâ die loco non movebatur, tacta bardè admodùm volitans quemlibet in locum impingebat, ova parit flava.

XXII. *Rosæ multiplex, media.*

Suprà Rosæ calycis nervulum inferius erucula flava residet, mense Julio in aureliam ex viridi rubram mutabatur, diebus quatuordecim elapsis elegans animalculum cinereum inde prodibat, ochræ colore lucido, albisque simul striis exornatum.

Similis eruca, exanimis adinstar, prosternebatur, effluxis

ali-

aliquot diebus ex eâ quinque erepebant acari , in doliola alba transeuntes , hæc autem memorata doliola contexebat eruca , periitque ; post dies quatuordecim quovis è doliolo parva musca prorupit.

XXIII. *Cerasus major fructu subdulci.*

PEr elegantis hujus erucæ dorsum nigra gemmis quasi intertexta vincula obliquè ferebantur , hisce foliis eam enutrivit , aliquandò deficiente pabulo , sibimet ipsis invicem esca fuère ; mense Julio ovalis figuræ telam argenti more fulgidam conficientes in aurelias immutabantur , Augusto phalenæ elegantes colore nigro , albo , cinereo , auriaco , roseoque colore præditæ , prodibant.

Supra cerasum erucula formosa celerrimi gressûs exhibetur , in Augusti principio fuscam in aureliam immutata , tandem in fine mensis ejusdem papilio , facta est , qualem figura repræsentat.

XXIV. *Rosa maxima multiplex.*

Hocce erucularum viride genus nigris constans capitibus , pediculos , calycesque , eodem , quo in inferiore rosæ calyce conspicis , modo , internè exedit , tactæ filo descendunt , eodemque mediante reascendunt , in fine Maii in aurelias fuscas immutantur , elapso quatuordecim dierum spatio fulgurantes ut aurum , volatu celeres , turbati dexterrimè sub folia sese recipiendi , peritissimi papiliones inde prodeunt.

XXV. *Fruetus grossulariæ , Spinosæ.*

Ingens in folio , propemodùm erecta demonstratur eruca , tacta , parte corporis superiore huc illuc agitur , trunculis posterioribus ità firmiter adhæret , ut illæsa vix aut ne vix quidem exfolvi possit ; in finem usque Maii eam enutrivit , tunc in aureliam fuscam invertebatur , elapso quatuordecim dierum spatio , albi , fusci , maculisque variis aspersi papiliones inde prorumpabant , quales inferior figura exhibet.