

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0032

LOG Titel: XXV. Fructus gorssulariae fativae, spinosae

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

aliquot diebus ex eâ quinque erepebant acari , in doliola alba transeuntes , hæc autem memorata doliola contexebat eruca , periitque ; post dies quatuordecim quovis è doliolo parva musca prorupit.

XXIII. *Cerasus major fructu subdulci.*

PEr elegantis hujus erucæ dorsum nigra gemmis quasi intertexta vincula obliquè ferebantur , hisce foliis eam enutrivit , aliquandò deficiente pabulo , sibimet ipsis invicem esca fuère ; mense Julio ovalis figuræ telam argenti more fulgidam conficientes in aurelias immutabantur , Augusto phalenæ elegantes colore nigro , albo , cinereo , auriaco , roseoque colore præditæ , prodibant.

Supra cerasum erucula formosa celerrimi gressûs exhibetur , in Augusti principio fuscam in aureliam immutata , tandem in fine mensis ejusdem papilio , facta est , qualem figura repræsentat.

XXIV. *Rosa maxima multiplex.*

Hocce erucularum viride genus nigris constans capitibus , pediculos , calycesque , eodem , quo in inferiore rosæ calyce conspicis , modo , internè exedit , tactæ filo descendunt , eodemque mediante reascendunt , in fine Maii in aurelias fuscas immutantur , elapso quatuordecim dierum spatio fulgurantes ut aurum , volatu celeres , turbati dexterrimè sub folia sese recipiendi , peritissimi papiliones inde prodeunt.

XXV. *Fruetus grossulariæ , Spinosæ.*

Ingens in folio , propemodùm erecta demonstratur eruca , tacta , parte corporis superiore huc illuc agitur , trunculis posterioribus ità firmiter adhæret , ut illæsa vix aut ne vix quidem exsolvi possit ; in finem usque Maii eam enutrivit , tunc in aureliam fuscam invertebatur , elapso quatuordecim dierum spatio , albi , fusci , maculisque variis aspersi papiliones inde prorumpabant , quales inferior figura exhibet.

Fructui erucula infidet in medio viridis, in parte verò corporis posteriore, anterioreque flavis, nigrisque punctis distincta; fructui huic non mediocriter damnum infert, repetitis vicibus exuvias deponit, venustiori continuo metamorphose elegantior facta sub finem Maii degenerat in nympham, trium septimanarum spatio muscæ flavæ prodeunt, uti latus habet.

XXVI. *Urtica urens major.*

HÆc herba nigricantis erucæ pabulum est, quasdam earum in scatulâ reservavi, quæ decimo quarto die Junii operculo adhærebant, in aureliis expallido flavas inversarum unam aperui curiositate stimolata, figuræ è folio pendentis inferiùs correspondebat, vigesimo octavo die ejusdem mensis papilio inde prodibat, cujus latus exteriùs fuscum, interiùs autem ex obscuro auriaci coloris erat, nigro purpureoque mirificè exornatum. Ex aureliarum maximâ musca nigra graveolens exibat, qualis, suprà aureliam folio insidens est depicta.

Acaros etiam in excrementis harum erucarum deprehendi in doliola nigra mutatos, in quatordecim dierum spatio, quales exhibentur, in lucem muscæ prodibant.

XXVII. *Salix acuto folio.*

IN pendulo longiore folio animalculum flavum punctis nigris intertextum exhibetur, decimâ quartâ die Junii folio adhærebat, & in nympham degeneravit, præterlapsis quindecim diebus, prodibat inde scarabæus, qualis folio superiùs infidet, qui semen suum adhuc in hæc folia disseminat, ex quo alia rursum animalcula producuntur.

Folium inferius eruculam fuscam exhibet, hisce foliis emanant rivi, sextâ die Junii telam confecit, & in eâ, in sequentis anni mensẽ usque Majum permanfit, tunc talis inde musca prodit, qualis infrâ depingitur.

XXVIII. *Rosa versicolor.*

Spadicea rosæ insidens eruca ventre suo graditur tribus solummodò trunculis anteriùs instructa, hisce rosis in finem usque