

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0063

LOG Titel: VI. Rosa centifolia, Batavica, rubra

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

V. *Lapathum acutum crispans.*

Super hac herbâ è rufo furvis lineolis in crucem sparfis variegata eruca reperitur, ab antè sex ungviculis, & à retrò sex pediculis instructa est, Majo in aureliam fuscam immutatur, undè fuscis lineis signata phalænula educitur, qualis superius volitans exhibetur.

Eâdem herbâ similem eruculam enutrivi, qualis ex alio latere tortuosa figuratur, è fusco psittacorum more virentibus lineis distinguebatur, pedetentim tamen flavescens tandem fusca facta est, gradu iners erat, sex ab antè ungviculis, à retrò quatuor adhuc pediculis constabat, iter promovens, posteriores prioribus pedes jungebat, corpore gibbum formans, ac ità progrediebatur; Majo vertebatur in aureliam è lucido fuscâ, cujus terminus superior capitis aviculæ formam refert, elapso quatuordecim dierum curriculo albicans indè phalænula prodiit rubris constans oculis, per cujus quatuor alas rubicunda linea cum margine pariter rubrâ ferebatur, in volando fuit celerrima.

VI. *Rosa centifolia rubra.*

FLos hicce formosus è lucido virentis erucæ pabulum est, albæ corpus ejus in longum decurrunt lineæ, quodlibet membrum puncto albicante, ac inferius lineâ flavâ est adornatum, sex ab ante ungviculos, in medio pediculos octo, & duos à retrò similes habet, Junio virentibus rosarum foliis sese intexit, ac in aureliam fuscâ immutatur, Julio mense, qualis superius exhibetur, educta fuit phalænula è lucido flavâ, cujus alæ inferiores obliquè sub aspectum cadentes auri more coruscant.

In Rosariorum etiam fronde simile vermium genus invenitur, qualis in rosæ pediculo vermis conspicitur; hicce Majo junioque mensibus vitâ fruitur, acarum colore refert, per dorsum rosei coloris lineâ adornatus, parvulis continuò circumvolantibus muscis tanquam pabulo vescitur, neque loco movetur, at per corpus ejus decurrentes attingere valens, probosci-

dem singulari dexteritate protrudit, & hoc animalculum captivum deprædatur, cujus humorem exfugit, reliquum verò ficum rejicit, tandem mutationem subit, & in bullam claram transfiguratur, diebus quatuordecim elapsis, qualis in rosæ calyce remonstratur, musca prorumpit, quæ alulas intricatas, atque convolutas, per medium circiter horæ spatium pediculis suis extricans, ac explicans avolat, capite constat flavo, rubris ambobus oculis, corpore superius existente viridi, inferius flavo nigroque colore tincto, sex fundata pediculis flavis; facili negotio, commodèque capitur.

VII. *Rosa sylvestris major.*

IN hisce rosis hujusmodi è lucido virescentes eruculæ repriuntur, ab antè sex ungviculis, & à retrò quatuor pediculis instructæ, in fine Maii fuscas in aurelias immutantur, undè quatuordecim diebus elapsis fimiles è lucido fuscæ phalænulæ producuntur, quales suprâ videre est. Infrâ hanc, punctulis viridibus intertexta, folium viride perrepat erucula, nigro capite, in medio mensis Maii in aureliam è lucido fuscam immutatur, in Junii principio prodit indè phalænula, cum alæ superiores, ungviculi que sex lineis flavis fuscisque variegatæ sunt, oculos habet nigros, & inferiores alæ cinereo tinctæ sunt colore. Ex alio latere viridis pariter hæret euca capite nigro, sex ungviculis ab antè, in medio totidem, & à retrò duobus adhuc pediculis constans, hanc solis, in finem usque Maii, rosis enutrivî, ubi in sese convoluta jacebat immota; elapsis aliquot diebus acarus inde candidus educebatur, qui in doliolum è lucido fuscum, ut in mediâ rosâ demonstratur, immutatus, in cœruleam exilem, elapso quatuordecim dierum curriculo, muscam transit, qualis paulò altiore loco apponitur.

VIII. *Galiopsis florens.*

HUjusmodi hirsutam nigricantem, flavis, albicantibusque punctulis exornatam erucam aliquo tempore hac herbâ enutrivî, sex ab antè ungviculis, in medio octo, & à retrò duo-