

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0071
LOG Titel: XIV. Meliotus lutea
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

in aureliam è lucido fuscā immutatur , quæ tacta fortiter mouetur , inde denique prodit ex albo fusca phalænula , qualis suprà volitans exhibetur.

Inferiùs ad dextram erucula fusca medio folio insidens conspicitur , quolibet in membro albis munita pediculis , hæc florē hujus arboris unā cum folio depascit , residet & in arboribus cydoneis , mense Majo candidam resplendentemque texturam conficit , & in aureliam immutatur , quatuordecim diebus elapsis , qualis superius unā repræsentatur , musca nigricans indè prodit.

Arborem albis variegata lineis , eruca viridis perreptat , cuius flos hicce pabulum est , cinerea admodūm est ; deponens exuvias roseum colorem induit , Augusto texturam parat albam , ac in doliolum fuscum immutatur , ita figurata permanet in annum usque sequentem , Julio musca nigra indè producitur , rubros habens oculos , qualis ad sinistram unā exhibetur.

XIII. *Lappa major.*

ETiamsi valdè sit amara hæc herba , dorsum tamen fuscis. Atrisque maculis conspersum habentis , subtūs flavæ mulcæ pabulum est , ab antè sex ungviculis , in medio octo pediculis constat , Junio mense in aureliam hepatici coloris immutatur , Julio phalæna indè prodit , capite , alisque superioribus hepatico pariter colore tinctis , cum fuscis auratisque punctulis , constans , corpus unā cum alis inferioribus flavum , aureumque colorem refert , sex ungvibus fuscis munitum , vespere volitat.

XIV. *Melilotus lutea.*

INferiores hæ erucæ prægrandes hac herbâ vescuntur , viridi quodam pallido maculis fuscis eleganter signato colore præditæ sex ab antè constant ungviculis , octo in medio , & a retrò duobus adhuc pedibus , quodlibet membrum corporis inferioris maculâ sanguineâ conspersum est ; tactæ turbantur admodūm , in fine Maii texturam conficiunt albicantem , &

in aurelias è lucido lividas immutantur, in medio Junii formosæ indè phalænulæ prodeunt, capite, corpore, ungvibus, cornulis, alisque superioribus cinereo colore tinctis, nigrisque ac candidis punctulis lineolisque conspersis, alis posterioribus elegantissimâ minii rubedine superbientibus constantes, quarum una folio insidens figuratur.

Herbeam hujuscemdi erucam superius depictam in Salice mense Julio deprehendi, eamque trifolio enutrivi in finem usque dicti mensis, ubi in aureliam janthinam immutabatur, mense Mayo sequentis anni è lucido purpureis lineis virgata phalænula exilis quidem, volatu tamen celerrima indè producta fuit.

XV. *Viola Lutea pleno flore.*

E Jusmodi ex obscurò cinereas per utrumque latus albicante condecoratas lineâ hirsutas erucas inferius in folio viridi exhibitas hoc super flore est reperire, easdem etiam suprà grossulariæ baccas, & halicacabos invenies, caput flavum, ut & antici sex ungviculi, & octò medii posticique duo pediculi cinerei sunt, Augusto mense in hepatici coloris aurelias ab aliâ parte, viridi super folio annotatas immutantur, hac interim figurâ signatæ permanent in Aprilem usque sequentis anni, ubi superius suprà florem depictæ phalænula capite, corpore, alisque è lucido flavis, maculis nigris exornatis, oculis verò, cornulis, ungviculisque nigris constantes, vespere volitantes educuntur.

Superius ab altera parte super flore è lucido virescens cum punctis flavis intertextis eracula remonstratur, hanc flore hoc in medium usque Junium enutrivi, ubi telam parabat albicantem, & in aureliam subviridem mutabatur, in fine dicti mensis alba maculis cinereis exornata phalænula prodiit cinereis duobus cornulis instructa, quatuorque ungviculis constans.

XVI.