

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0074

LOG Titel: XVII. Carpinus florens

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XVI. *Flos cuculi, & gramen vulgare.*

HOsce flores in complementum ac ornamentum hic ad-
 junxi, sed hæc subrepens prægrandis eruca fuscis macu-
 lis aspersa, flavis, albisque punctulis variegata, capite pilo-
 rum nigrorum fasciculum, à tergo aculeum instar cornuli, ac
 ab utroque latere curiosos pilorum alborum pariter fasciculos
 gestitans, pilorum inquam tantoperè dependentium, ut ne
 pedes quidem in conspectum venire possint, in gramine reperitur,
 quod avidissimè depascitur; humido valde delectatur, in me-
 dio mensis Maii subflavam parat texturam, ac in aureliam fla-
 vam immutatur, in fine prædicti mensis venit in conspectum
 phalæna, qualis suprà cernitur, flava, fusca superiores alas li-
 nea transversa decurrit, quæ duabus etiam utrinque maculis,
 cum marginibus fuscis, notatæ sunt; alba parit ovula.

Animadverti etiam, has erucas humi silentiosè sterni, & ex
 unâquâque erucâ quatuor magnos paulò post acaros exire, qui
 totidem in doliola mutati, tot etiam in ingentes muscas im-
 mutati fuerunt.

XVII. *Carpinus florens.*

IN pede figuræ flava lineis, punctisque nigricantibus distinc-
 ta eruca demonstratur, sex ab antè nigris ungviculis, in
 medio octo, & à retrò duobus adhuc flavis pedibus constans,
 tarda est admodum, tacta aliarum more erucarum immota
 jacet, in finem usque Junii hocce folio illam enutrivit, ubi tex-
 turam conficiebat albam, chartæ pergamenæ tenuiori simi-
 lem, ac argenti more fulgidam, in hac fuscam in aureliam im-
 mutata est, in medio sequentis mensis Julii phalænula Janthi-
 na inde prodiit, cujus superiores alæ maculis rubicundis per
 obliquum adornatæ mirificè coruscabant, inferiores alæ pari-
 ter rubicundis maculis aspersæ erant, in volando segnes sunt
 hæ phalænulæ, floribus in gramine unâ communiter insident,
 flava pariunt ovula.

Ad dextram etiam herbea folio insidet erucula, in prunis
 quoque reperibilis ab antè sex ungviculis, in medio octo & à

retro duobus adhuc pediculis constat, tacta multum Agitur, Julio mense in aureliam fuscam immutatur, decursu quatuordecim dierum phalænula capite, cornulis, pediculis, & alis superioribus hepatico, cum lineolis fuscis, inferioribus autem alis cinereo, colore tinctis constans indè prodiit, humilis in gramine volatus ejus est.

XVIII. *Milii folium terrestre florens.*

HOcce flori superius insidens animalculum in acetosâ quoque reperitur, nigri coloris est, sex constans ungviculis, sub finem Maii flavam in aureliam immutatur capiti canino non absimilem, elapso quatuordecim dierum intervallo, scarabæolus indè niger erumpit, qualis humi currens exhibetur.

Ex alterâ parte animalculum exile reperit humi, quod in gramine reperi, ejusdem cum pediculis, coloris est, in fine Junii scatulæ operculo firmiter adhæret, in quâ illud enutriebam, convoivebatur aliquantulum, fuscum, duriusculumque factum, elapsis duodecim diebus indè scarabæolus ruber, punctisque nigris aspersus exiit, qualis unâ humi exhibetur.

XIX. *Flos tinctorius.*

INferius ad dextram viridi folio viridis insidet eruca, cujus per dorsum recta nigricans, inferius autem ab utrâque parte alba decurrit linea, tacta jacet aliquot horis immobilis, in medio Junii texturam quadrangulari modo perforatam, piscium reti similem, conficiebat, tacta violento motu agitur aurelia, sub initium Julii è lucido viridis, albas & ex atro virides obliquè per alas decurrentes lineas gestans, inde phalænula prodiit, in volando celerrima.

Ad sinistram cerasi folium humi jacet, cui nigricans insidet eruca qualibet in incisione superius cærulea duo, fasciculo pilorum exornata coralliola gestans, hujus, cum capite, ungviculi sex, octo medii, duoque postici pediculi flavo ochræ colore tincti sunt; in fine Junii cinereâ sibi texturâ paratâ fuscâ in aureliam immutatur nigris in capite pilis obsitam, in medio Julii fulvâ lineis fuscis virgata prodiit indè phalæna,