

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0079
LOG Titel: XXII. Flos mali
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XXII. *Flos mali.*

Pilosam hanc & flavam lineis obliquis rotundis exornatam erucam in malo deprehendi, caput habebat fuscum, ac retrò illud cœrulea duo, ac quolibet in membro duo rosea coralliola, eodem colore tintæ fuere pediculi, utrumque latus fusci pilorum fasciculis protuberavit, in Julii principio deponebat exuvias, telam perforatam texebat flavam piscium reti similem, & in aureliam hepatici coloris immutabatur, per totum corpus pilorum fasciculis obsitam, ac inferius in apice nodulo signatam, tacta tam violento motu agitabatur, ut diversis vicibus sursum deorsumque volveretur; quatuordecim diebus elapsis inde nigricans albicansque phalænula prodiit, rubicunda duo cornula lata protendens, pennigera, & retrò caput inter cornula rubram gestans lineam, celer erat, vespere volitans.

XXIII. *Vitis florens.*

Ambæ prægrandes hæ erucæ sunt ejusdem generis, colore solummodo differentes, superior enim fulvo, sed inferior viridi colore tincta est, utraque punctis lineisque nigris & albis exornatur, de vite pabulum sumunt, pascentes, quiescentibus tertiaro longiores amplè extenduntur, harum excrementa ex obscuro viridia quinque formant angulos, anguli tamen rotundi bacellulos in fasciculum compositos repræsentant, quemadmodum pro facilitiore intelligentiâ inferiori erucæ appicta conspicis.

Ex animadverso jam habui, viridum erucarum aliquas humi Stratas, in sece compendiosissimè convolutas fuisse, unde paulò post sex acari prodierunt, hi mutabantur in fusca do-liola, exinde vero ex obscuro cœruleæ lineis nigris virgatae muscæ fuerunt eductæ, oculos habentes rubicundos, alasque vitri more perspicuas.

Eruca fusca in medio Julii fuscam in aureliam immutabatur ac itâ transfigurata in sequentis anni mensem usque majum permanxit, ubi formosa phalæna prodiit, cuius caput, corpus, ac alæ superiores eleganti roseo colore tintæ, lineis maculisque, psit-