

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0086
LOG Titel: XXIX. Gallium luteum
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XXVIII. *Absynthium vulgare.*

Quantumvis etiam acerba ac amara herba qualiscunque fucrit, certo tamen animalium generi semper in pabulum cedit, quod in hoc stipite nobis est experientiae; hæcce stipiti infidens eruca albicante linea maculisque fuscis per dorsum signata, per latera è gilvo viridi, cum lineis fuscis, colore distincta, inferius albicante, quolibet in membro notata puncto, ab ante sex ungviculis in medio octo, & à retro duobus adhuc pediculis instructa, caput huc illuc, contrectata, celeriter agitat: mense Majo nigram in aureliam immutatur, elapo trium hebdomadarum spatio producitur inde phalænula, cuius caput, alæque superiores subrufæ nigris, fuscisque lineis, punctisque condecoratae sunt, in medio quoque superiorum alarum albicans argenti more splendida macula conspicitur, inferiores alæ cum corpore fuscae sunt, cornula nigra duo gestat, oculos duos habet coruscantes, vespere volitat.

Inferius erucula subviridis, longiuscula cernitur, curiosè incedens, partem corporis posticam anticæ jungit, monstrosum formans gibbum, ac ita suum iter prosequitur, per vices aliquamdiu suis erecta stat pedibus posterioribus.

In fine Julii viridem in aureliam immutatur, elapo quatuordecim dierum spatio producitur inde phalænula, cuius caput, corpus, alæque superiores albis, nigris, fuscisque lineolis ac punctis signatae, inferiores autem è lucido fuscae sunt: duo quoquè gestat è lucido fusca cornula, quatuor nigris punctis variatos ungviculos habens, in volando est celerrima.

XXIX. *Gallium luteum.*

HÆcce herba formosæ, viridis erucæ pabulum est, cuius à parte corporis superiore rotunda multa, coloris albi, exaltata punctula, in utroque latere alba linea conspicitur, à tergo cornu semi-albo semique-cærulæ, munita, sex ab ante ungviculis, in medio octo, & à retro duobus abhuc rosei coloris pediculis instructa est; in medium usque Julium eam enutrivi, ubi pelle suâ exuebatur, roseum induens colorem in

flavam subcinereum tandem aureliam immutabatur, hocce modo transformata dum attingebatur impatiens admodum erat. In principio Augusti phalænula indè prodiit, cuius caput, pars corporis antica, unguiculi, alæque superiores fusci coloris cum lineis obscuris erant, corporis autem pars postica albicans lineis nigris transversis lanugineque decora, alæ inferiores auraci coloris erant, oculis nigris constabat, nigrisque corniculis, ac proboscide munita erat, ad alimentum è floribus extra-hendum, de die volitat, alacris est, captuque difficillima. Acarum albicanter, capite nigro hic adjunxi, hunc in panno deprehendi, in quo ovalis figuræ domunculam laneam confercerat, licet perforatam, uno termino contrectato, per alium terminum celeriter exiens iteratò redibat, postmodum in vasculum immutatus post quaturdecim dierum spatum in phalænulam transit, argenti more splendidam, lineolis nigris punctulisque similibus exornatam, cum tamen à damno quod infert, abundè innotuerit, hisce jam dictis acquiesco.

XXX. *Alni folia.*

HÆc folio irreps eruca lineis albis nigrisque, similibus que punctis variegata, nigro constat capite, sex ab ante unguiculis, in medio duodecim, &c à retro duobus adhuc pedibus, in finem usque Junii hisce foliis eam enutrivi, ubi folio involuta texturam conficiebat albicanter, ac in aureliam immutabatur, elapsò quatuordecim dierum spatio alba punctis nigris signata, qualis superius exhibetur, inde phalænula prodiit. Folio infimo nigrum punctis flavescientibus notatum turpe animalculum irrepit, sex unguiculis instructum, hoc in medium usque Junitum hisce foliis enutritum petiolo firmiter adhærebat, ac exutâ priore formâ, nigrescebat, albisque maculis variegabatur, paucis diebus effluxis duplex scarabæolorum genus, utrumque nigrum atque flavum, signaturis tamen distinctum aliis atque aliis, indè prodiit.

Infrà propè petiolum proreperit erucula flava, quam in finem usque Junii hisce foliis pariter enutrivi, ex cuius adhuc vivo corpore tres acari tunc erepebant, hi albicanibus ovulis fere intexuerunt, ac in doliola sunt immutati, unde post duodecim dies