

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0087

LOG Titel: XXX. Alni folia

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

flavam subcineream tandem aureliam immutabatur, hocce modo transformata dum attingebatur impatiens admodum erat. In principio Augusti phalænula indè prodiit, cujus caput, pars corporis antica, ungviculi, alæque superiores fusci coloris cum lineis obscuris erant, corporis autem pars postica albicans lineis nigris transversis lanugineque decora, alæ inferiores auriaci coloris erant, oculis nigris constabat, nigrisque corniculis, ac proboscide munita erat, ad alimentum è floribus extrahendum, de die volitat, alacris est, captuque difficillima. Acarum albicantem, capite nigro hic adjunxi, hunc in panno deprehendi, in quo ovalis figuræ domunculam laneam confecerat, licet perforatam, uno termino contrectato, per alium terminum celeriter exiens iteratò redibat, postmodum in vasculum immutatus post quatuordecim dierum spatium in phalænulam transit, argenti more splendidam, lineolis nigris punctulisque similibus exornatam, cum tamen à damno quod infert, abundè innotuerit, hisce jam dictis acquiesco.

XXX. *Alni folia.*

HÆcce folio irrepens eruca lineis albis nigrisque, similibusque punctis variegata, nigro constat capite, sex ab ante ungviculis, in medio duodecim, & à retro duobus adhuc pedibus, in finem usque Junii hisce foliis eam enutrivit, ubi folio involuta texturam conficiebat albicantem, ac in aureliam immutabatur, elapso quatuordecim dierum spatio alba punctis nigris signata, qualis superius exhibetur, inde phalænula prodiit. Folio infimo nigrum punctis flavescens notatum turpe animalculum irrepit, sex ungviculis instructum, hoc in medium usque Junium hisce foliis enutritum petiolo firmiter adhærebat, ac exutâ priore formâ, nigrescebat, albisque maculis variegabatur, paucis diebus effluxis duplex scarabæolorum genus, utrumque nigrum atque flavum, signaturis tamen distinctum aliis atque aliis, indè prodiit.

Infra propè petiolum prærepat erucula flava, quam in finem usque Junii hisce foliis pariter enutrivit, ex cujus adhuc vivo corpore tres acari tunc erepebant, hi albicantibus ovulis sese intexuerunt, ac in doliola sunt immutati, unde post duodecim dies

dies tres in lucem muscæ proruperunt, nigro capite quælibet harum, ac corpore ungviculisque flavis constabat.

Inferius ad sinistram cinerea punctisque nigris notata adhuc proreptat erucula, capite ungviculisque anterioribus nigris, cæterisque pediculis albicantibus constans, hæcce Junio mense velut exanimis humi sternebatur, elapsis aliquot diebus oblongus tenuis, è corpore vermibus flavus prodibat instar fili chalybæi, contemplatione dignus erat hicce vermibus, variis enim flexuris contortus ac implicitus extingvebatur.

XXXI. *Cariophyllus variegatus.*

FLori huic primò florescenti per dorsum ex obscuro fusca pedibus verò è lucido flavis constans eruca insidet, hocce flore in medium usque Julium per me enutrita fuscam in aureliam immutabatur, taliter figurata in medium usque sequentis anni Junium permanfit, ubi tandem, capite, alisque superioribus nigris, corpore postico, alis inferioribus, cornulis, ungviculisque cinereo colore tinctis constans phalæna prodiit, vespere volitabat.

Vermiculus alius in majoris floris petiolo exhibitus, viridibus, hoc flore pariter enutritis, animalculis pascitur; dictus vermibus neque loco, neque corpore movetur, ast antè memoratos pediculos virescentes per corpus decurrentes avidè celeriterqueprehendit, omnemque fuccum, folliculo vacuo rejecto extrahit; sub finem Junii in bullam immutabatur, unde effluxo quatuordecim dierum curriculo musca prodiit, cujus caput rubicundum, corpus nigrum, lineis transversis flavis variegatum, pediculi que fuscæ fuerant.

XXXII. *Borago cærulea.*

Sub dextrâ folio viridis insidet eruca, ab antè sex ungviculis, & à retro quatuor pediculis constans, super brassicâ cypriâ sive cauliflorâ pariter reperitur, Augusto mense telam conficit albicantem, harum erucarum aliæ in fuscas, in aurelias ex obscuro virides aliæ immutantur, aliæ inter texendum contortæ, & convolutæ transeunt in acaros, qui subito exting-