

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0096
LOG Titel: XXXIX. Brassica viridis
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XXXVIII. *Eryngium Campestre.*

Simile nigricantium erucarum genus, quarum una inferiori folio insidet, in urticis invenitur; cum in primâ parte jam delineaverim urticam, eringium hoc campestre pulchritudinis suæ cauſâ hic appingere volui.

Sub finem Junii metamorphosin subeunt hæ erucæ, à retrò fixæ deorsum propendent, evadentes fuscæ velut auro coruscantes aureliæ, spatio quatuordecim dierum elapso formosi papiliones inde prodierunt, quorum alæ inferiores ferruginei, superiores autem paulò lucidioris erant coloris, maculis albicantibus aspersæ, corpore nigro, cornulis ex albo aculeatis, unguiculis cinerei coloris constabant, in volando celeres.

Sedet & infrà viridis erucula capite flavo, hæc in urticis etiam invenitur, quarum folia convolvit, & in ea se recipit, in eisdem mutatur in aureliam fuscam, elapso quatuordecim dierum spatio exilis è lucido fusca lineis nigris virgata phalænula producitur; harum erucarum aliquæ prostratæ emoriuntur, è quibus parvuli quâm plurimi prodeunt Acari, brevi temporis spatio mutantur in doliola, unde post decem aut duodecim dies parvulæ muscæ nigricantes exeunt.

XXXIX. *Brassica viridis.*

Plantam hanc recentem adhuc, herbea eruca gulosè depascit, quæ flavâ per dorsum lineâ, & in utroque membro cuiuslibet latere punctulis nigris signata est, perfectam noctâ magnitudinem pelle suâ sese exuit, alibi suspenditur, ac in fuscum punctis nigris notatam aureliam immutatur, elapso quatuordecim dierum spatio papilionum priùs demonstratis minorum genus indè educitur.

Augusto mense eruculas in brassica quoque deprehendi, vi-flava per dorsum lineâ signatas, quolibet in membro duobus pediculis instructas, hæ in aurelias erucæ parùm absimiles immutabantur, elipsis decem diebus muscæ, corpore nigris albisque maculis asperso, rubris oculis, unguiculisque sex flavis constantes in volandæ tardæ indè prodierunt.