

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis
Autor: Merian, Maria Sibylla
Verlag: Oosterwyk
Ort: Amstelædami
Jahr: 1717
Kollektion: Zoologica
Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen
Werk Id: PPN476830389
PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>
OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0098
LOG Titel: XLI. Urticae urentis folia
LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

XL. *Acetosa pratensis florens.*

Super hac herbâ nigricans rubris albisque maculis aspersa lagninosa deprehenditur eruca, cuius pili cum capite, sex anticis ungviculis, & octo pediculis è lucido fuscâ sunt, perfectam confecuta magnitudinem aliò cum flexurâ sese collocat, lucescit, & in aureliam è lucido fuscâ immutatur, aliquæ tales trium mensium spatio, aliæ in subsequentis anni mensem Aprilem usque permanebant; ubi tandem exilis inde phalænula prodiit, cuius corpus alulæque inferiores flavæ, caput cum alulis superioribus, cornulis, ungviculisque fuscum aculeis nigris instructum erat, vespere volitat.

XLI. *Urticæ urentis folia.*

TRes hic erucæ generis quidem ejusdem, diversi tamen coloris exhibentur, superior viridi folio insidens in totum nigricans punctulis albis fulgentibus supra pilorum radicem, ac inferiùs ab utroque latere linea flavâ est exornata, in medium usque Septembrem per me enutrita sursum deorsum propendens fuscâ in aureliam immutabatur, elapso quatuordecim dierum spatio formâ alias atque specie longo post se intervallo relinquens indè papilio prodiit, constans corpore, capite, cornulis, ungviculisque ferruginei coloris, latâ minii rubedinem referente lineâ, cæruleis albisque maculis signatus, alulæ inferiores haud vulgari pulchritudine sunt donatæ, diversis è variorum colorum invicem mixtorum genere ornamentulis illustratæ.

Sub hac ad dextram collocata eruca corpore è lucido flavo, capite fusco, ungviculis pediculisque roseo colore tinetis constat, simili metamorphose fuscâ in aureliam immutabatur, unde pariter elapso quatuordecim dierum curriculo superiùs allegato similis papilio productus fuit.

Tertia in rotundum flexa eruca priùs nigra colorem immutavit, fusca facta, & humi sternebatur in rotundum flexa, ex quâ brevi temporis spatio quamplurimi exiles acari prorepebant, qui in unum sese contexentes in doliola tandem immutabantur,

duodecim diebus elapsis, qualis hic una exhibetur nigra quolibet è doliolo musca prodiit.

Infimo folio fusca pilosaque erucula insidet, cuius cum ventre dorsum rubescet, tacta convolvebatur instar globuli, sub finem Maii humi flexa sternitur, ac in medium usque junium immota jacet, ubi tandem musca prodit, cuius caput, corpus & cornula nigro, quatuor alulae pellucido, ungviculique flavo colore tincti erant, in postica parte maculâ nigrâ conspersa fuit.

XLII. Linum florens.

Hicce flos erucæ subviridi flavæ cedit in pabulum hæc inferiùs è lucido viridi colore tincta ab utroque latere albicante lineâ signata est, ab ante sex ungviculis, in medio octo, & à retrò duobus adhuc pediculis constat, in Augusti principio depositis exuviis coruscantem in aureliam immutabatur, ultimâ subsequentis anni mensis Maii die cinerea prodiit indè phalæna, cuius alæ nigro colore erant exornatae, nigris constabat oculis, sub vesperam volitabat.

XLIII. Centumnodia, sive polygonum florens.

Ad sinistram fusca collocatur eruca, inferiùs cinerea, ab antè sex ungviculis, in medio octo, & à retro duobus adhuc pediculis constabat, sub finem Julii in aureliam è lucido fuscum immutabatur, in medio Augusti cinerea lineis nigris virgata indè prodiit phalænula vespere volitans. Inferiùs adhuc ad dextram eruca viridis exhibetur, tribus flavis per corpus lineis, pluribusque maculis nigricantibus exornata, hac herbâ in finem usque Julii ipsam enutrii, ubi jacebat immota, elapsa quatuordecim dierum spatio parvi triginta quatuor albicantes acari exinde prorepebant, exiguo temporis spatio in doliola mutati, ex quibus sub finem Julii parvæ totidem muscae nigræ proruperunt.

Alia pari modo hujus generis eruca humi silentiosè sternebatur, unde magnus albicans vermis prorepsit, hic in doliolum fuscum immutatus in ingentem fuscum lineis nigris variegatam muf-