

Werk

Titel: Erucarum ortus, alimentum et paradoxa metamorphosis

Autor: Merian, Maria Sibylla

Verlag: Oosterwyk

Ort: Amstelædami

Jahr: 1717

Kollektion: Zoologica

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN476830389

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN476830389>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=476830389>

LOG Id: LOG_0099

LOG Titel: XLII. Linum florens

LOG Typ: chapter

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

duodecim diebus elapsis, qualis hic unâ exhibetur nigra quolibet è doliolo musca prodiit.

Infimo folio fusca pilosaque erucula insidet, cujus cum ventre dorsum rubescebat, tacta convolvebatur instar globuli, sub finem Maii humi flexa sternitur, ac in medium usque junium immota jacet, ubi tandem musca prodit, cujus caput, corpus & cornula nigro, quatuor alulæ pellucido, ungviculi-que flavo colore tincti erant, in postica parte maculâ nigrâ conspersa fuit.

XLII. *Linum florens.*

Hicce flos erucæ subviridi flavæ cedit in pabulum hæc inferiùs è lucido viridi colore tincta ab utroque latere albicante lineâ signata est, ab ante sex ungviculis, in medio octo, & à retrò duobus adhuc pediculis constat, in Augusti principio depositis exuviis coruscantem in aureliam immutabatur, ultimâ subsequenti anni mensis Maii die cinerea prodiit indè phalæna, cujus alæ nigro colore erant exornatæ, nigris constabat oculis, sub vesperam volitabat.

XLIII. *Centummodia, sive polygonum florens.*

AD sinistram fusca collocatur eruca, inferiùs cinerea, ab antè sex ungviculis, in medio octo, & à retrò duobus adhuc pediculis constabat, sub finem Julii in aureliam è lucido fuscâ immutabatur, in medio Augusti cinerea lineis nigris virgata indè prodiit phalænula vespere volitans. Inferiùs adhuc ad dextram eruca viridis exhibetur, tribus flavis per corpus lineis, pluribusque maculis nigricantibus exornata, hac herbâ in finem usque Julii ipsam enutrivit, ubi jacebat immota, elapso quatuordecim dierum spatio parvi triginta quatuor albicantes acari exindè prorepebant, exiguo temporis spatio in doliola mutati, ex quibus sub finem Julii parvæ totidem muscæ nigræ proruperunt.

Alia pari modo hujus generis eruca humi silentiosè sternebatur, unde magnus albicans vermis prorepsit, hic in doliolum fuscum immutatus in ingentem fuscâ lineis nigris variegatam mus-