

Werk

Jahr: 1984

Kollektion: fid.geo

Signatur: 8 Z NAT 2148:55

Digitalisiert: Niedersächsische Staats- und Universitätsbibliothek Göttingen

Werk Id: PPN1015067948_0055

PURL: http://resolver.sub.uni-goettingen.de/purl?PPN1015067948_0055

LOG Id: LOG_0019

LOG Titel: Long-term variations recorded by extensometers

LOG Typ: article

Übergeordnetes Werk

Werk Id: PPN1015067948

PURL: <http://resolver.sub.uni-goettingen.de/purl?PPN1015067948>

OPAC: <http://opac.sub.uni-goettingen.de/DB=1/PPN?PPN=1015067948>

Terms and Conditions

The Goettingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept the Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact

Niedersächsische Staats- und Universitätsbibliothek Göttingen
Georg-August-Universität Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
Email: gdz@sub.uni-goettingen.de

*Short communication***Long-term variations recorded by extensometers****P. Varga**

Eötvös Lorand Geophysical Institute of Hungary, Columbus u. 17–23, H-1145, Budapest, Hungary

Key words: Extensometer – Secular strain rates

On the basis of stress accumulation between two large earthquakes the maximum strain rate can be estimated. We suppose that the rigidity is $\mu = 3 \times 10^{10}$ N/m² and for the rock strength we take $p \leq 10^7$ N/m² (Latynina and Karmaleiva, 1978). If the time interval between two big earthquakes is $t = 100$ years, then the maximum strain rate is $p/(\mu t) \leq 3 \times 10^{-6}$ /year. This value could possibly be higher in tectonically very active regions (where t could also be smaller), but certainly not in quiet areas.

To compare this value with the recorded secular rates we compiled Table 1, where the mean secular rates are listed for 28 different stations distributed all over the world. Presently there are about 125 extensometric stations in operation (Bilham, 1973). The data presented in Table 1 can

be assumed to characterize the whole network. The records used usually cover a period of more than 1 year. (An exception is the series recorded by Berger and Levine (1974) in the Poorman mine with a laser strainmeter, which covered only 9 months.) The mean of the secular rate values is 2.1×10^{-6} /year. A rough estimation of the horizontal crustal movements resulting from this value is 2.1 mm/year in 1 km, a value which can be measured by geodetic techniques. For a Young's modulus of $(8-12) \times 10^{10}$ N/m² (Latynina and Karmaleiva, 1978) the corresponding stress rate is $(1.7-2.5) \times 10^5$ Nm⁻²/year.

There is a tendency in the literature to interpret such "secular strain rates" (determined with extensometers) tectonically. The question arises whether this is possible. We must remember that the values in Table 1 were determined at different times, with different types of instruments. The strainmeter used had different orientations and, what is more, in the case of stations where observations were carried out in different azimuths their mean value was used.

Table 1. Average secular strain rate values

Continent (Country)	Station	Type of extensometer	Secular rate of strain $\times 10^{-6}$ /year	Reference
<i>North America</i>				
USA	California	Q	1.0	Benioff, 1959
	Odensburg	Q	0.1	Beavan and Bilham, 1979
	Poorman mine	L	0.6	Berger and Levine, 1974
	La Jolla	L	0.2	Berger and Wyatt, 1973
	Nevada	Q	2.0	Priestley, 1974
	Pinon Flat	L	0.3	Wyatt and Agnew, 1983
	Aleuts	Q	3.0	Butler and Brown, 1972
	Mean		1.0	
<i>Europe</i>				
Great Britain	Yorkshire	W	0.2	Bilham et al., 1972
		L	0.7	Gouly et al., 1974
		W	5.0	Beavan, 1976
		W	5.0	Evans et al., 1979
GDR	Tiefenort	W	2.2	Harvardt and Simon, 1974
FRG	Schiltach	W	10.0	Kind and Emter, 1974
Hungary	Budapest	Q	2.0	This paper
USSR	Protvino	Q	0.5	Latynina, 1975
	Tbilisi	Q	4.0	Latynina and Karmaleiva, 1978
	Mean		3.2	

Table 1 (continued)

Continent (Country)	Station	Type of extensometer	Secular rate of strain $\times 10^{-6}$ /year	Reference
<i>Asia</i>				
Japan	Osakayama	W	2.0	Ozawa, 1971
	Amagase	W	1.2	Takada et al., 1981
	Kamigao	W	4.0	Nishimura et al., 1959
	Esashi	Q	0.2	Sato et al., 1983
	Erimo	W	0.2	Kasahara, 1981
USSR	Talgar	Q	1.0	Latynina, 1975
	Garm	Q	0.5	Latynina, 1975
	Turgen	Q	6.0	Ospanov et al., 1981
	Tschusal	Q	1.5	Latynina and Karmaleiva, 1978
	Inguri	Q	2.0	Latynina and Karmaleiva, 1978
			Mean	1.8
<i>Australia Oceania</i>				
Australia		Q	2.0	Sydenham, 1974
New Zealand		W	5.0	Gerard, 1973
		Mean	3.5	
		Mean of all measurements	2.1	

Q – quartz extensometer; W – wire extensometer; L – laser extensometer

In spite of these limitations we can conclude that secular rate values of the order of 10^{-6} /year cannot be simply connected with the degree of tectonic activity. The lowest values in Table 1 cannot be so easily discarded for tectonic interpretation, but the non-tectonic sources of the larger strain rates should be clearly understood before such an interpretation is made. Long-term variations at stations in Asia (all of which are in the tectonically most active areas) are similar to the ones observed in Europe, where the majority of the stations are in quiet regions. It can also be easily concluded that almost everywhere – regardless of the tectonic conditions – the long-term variations have values which are very close to the extreme rate expected.

It is necessary to add that the influence of external forces (meteorological, hydrological) on the long-term part of the strainmeter record – in the case of appropriate installation – are of secondary importance. According to the author's experience with a 21-m-long quartz tube extensometer installed in a 35-m-deep station in Budapest, the annual variations of the record are 10^2 times smaller than the secular rate. The influence of barometric and water table variations is even less.

It can be concluded that the strain rates cannot be used to estimate directly the magnitude of the tectonic processes. The estimates of stress rates and horizontal crustal movement values are bigger than the real values in nature.

Due to the significant differences among stations situated in the same region it seems that secular rates are modified or increased to a considerable extent by local conditions of different type, which are not connected to tectonic activity.

Acknowledgements. The author thanks the Alexander von Humboldt Foundation, which sponsored the work. The author is grateful to W. Zürn and D. Emter for their help and for critical discussions.

References

- Beavan, R.J.: Earth strain: ocean loading and analysis. Ph. D. Thesis, Cambridge University, 1976
- Beavan, R.J., Bilham, R.: Long series of strain observations from an aseismic area. EOS Trans. A.G.U. **60**, 316, 1979
- Benioff, M.: Fused-quartz extensometer for secular tidal and seismic strains. Bull. Geol. Soc. Am. **70**, 1019–1032, 1959
- Berger, J., Levine, J.: Spectrum of earth strain from 10^{-8} to 10^2 Hz. J. Geophys. Res. **79**, 1210–1214, 1974
- Berger, J., Wyatt, F.: Some observations of earth strain tides in California. Philos. Trans. R. Soc. Lond. A. **274**, 267–277, 1973
- Bilham, R.: The location of earth strain instrumentation. Philos. Trans. R. Soc. Lond. A. **274**, 239–241, 1973
- Bilham, R., Evans, R., King, G., Lawson, A., McKenzie, D.: Earth strain tides observed in Yorkshire, England with simple wire strainmeter. Geophys. J. Astron. Soc. **29**, 473–485, 1972
- Butler, D., Brown, P.L.: Secular strain measurements before and after the Caninikin nuclear test. Bull. Seismol. Soc. Am. **62**, 1455–1458, 1972
- Evans, R., Beavan, R.J., Bilham, R., King, G.: A survey of earth strain tides in Great Britain. Geophys. J. R. Astron. Soc. **57**, 119–135, 1979
- Gerard, V.B.: New Zealand Earth strain measurements. Phil. Trans. R. Soc. Lond. A. **274**, 311–321, 1973
- Gouly, N.R., King, G.C., Wallard, A.J.: Iodine stabilized laser strainmeter. Geophys. J. Astron. Soc. **38**, 377–387, 1974
- Harwardt, H., Simon, D.: Beobachtungen mit einem Drahtstrainmeter in der Station Tiefenort. In: 2. Internationales Symposium „Geodäsie und Physik der Erde“, Potsdam, Mai 1973, pp: 197–205 Veröff. ZIPE, Nr. 30, Teil I, 1974
- Kasahara, K.: Abnormal crustal strain change observed at Erimo before and after the Earthquake of the Nemuro Peninsula. J. Geodetic Soc. Jpn **27**, 282–288, 1981
- Kind, R., Emter, D.: Möglichkeiten zur Messung tektonischer Deformationen. Vortrag 34. Jahrestagung der Deutschen Geophysikalischen Gesellschaft, Berlin, 1974
- Latynina, L.A.: On horizontal deformations at faults recorded by extensometers. Tectonophysics **29**, 421–427, 1975

- Latynina, L.A., Karmaleiva, R.M.: Strain measurements, Moscow: Nauka (in Russian), 1978
- Nishimura, E., Kishimoto, J., Tanaka, I.: Observed secular tilting motion of the ground which preceded the occurrence of several destructive earthquakes. *Tellus* **11**, 15–22, 1959
- Ospanov, A.B., Latynina, L.A., Tihomirov, A.V.: Strain observation results obtained on the Prognostic Polygon Alma-Ata in connection with the seismicity. In: *Sovremennije dvizenija zemnoj kori*; pp: 21–24, Moscow, Nauka (in Russian), 1981
- Ozawa, I.: Observations on the crustal movement in Osakayama Tunnel of the former Tokaidao line. *J. Seismol. Soc. Jpn* **27**, 313–320, 1974
- Priestley, K.: Crustal strain measurements in Nevada. *Bull. Seismol. Soc. Am.* **64**, 1319–1328, 1974
- Sato, T., Ooe, M., Sato, N.: Tidal tilt and strain measurements and analyses of the Esashi earth tides station. In: *Proc. IXth Intern. Symp. Earth Tides*, 223–237, Stuttgart, E. Schweizerbart'sche Verlagsbuchhandlung, 1983
- Sydenham, P.H.: 2000 hr comparison of 10 m quartz-tube and quartz-catenary tidal strainmeters. *Geophys. J.R. Astron. Soc.* **38**, 377–387, 1974
- Takada, M., Furuzawa, T., Takemoto, S., Yamada, M.: Observation of crustal deformation at Amagase Observatory. *J. Geodetic. Soc. Jpn* **27**, 293–296, 1981
- Wyatt, F., Agnew, D.C.: High-precision deformation measurements in Southern California. In: *UGGI XVIII General Assembly*, p. 96, Hamburg Inter-disciplinary Symposia, Programme and abstracts, Vol. I, 1983

Received January 1, 1984; Revised version March 29, 1984

Accepted April 3, 1984